### PAPER • OPEN ACCESS

# He's polynomials method for analytical solutions of telegraph equation

To cite this article: S O Edeki et al 2019 IOP Conf. Ser.: Mater. Sci. Eng. 640 012117

View the article online for updates and enhancements.

## He's polynomials method for analytical solutions of telegraph equation

#### S O Edeki<sup>1</sup> O M Ofuyatan<sup>2</sup> and G O Akinlabi<sup>3</sup>

<sup>1,3</sup>Department of Mathematics, Covenant University, Ota, Nigeria <sup>2</sup>Department of Civil Engineering, Covenant University, Ota, Nigeria

Corresponding e-mail address: soedeki@yahoo.com

Abstract. In this paper, He's polynomials solution method (HPSM) is fully utilized for solving telegraph equation. The proposed HPSM is technically presented and applied to homogeneous linear form the telegraph equation. The results are expressed in closed form with good agreement compared to those in literature thereby attesting to the efficiency and reliability of the method as proposed. The HPSM remarked to be less time consuming with high level of accuracy. As such, it can serve as alternative to other methods.

Keywords: Analytical solutions; He's polynomials; telegraph equation, Adomian decomposition

#### 1. Introduction

In most physical and mathematical situations, modelling of physical phenomena leads to differential models in the form of equations which can be termed linear or nonlinear. Whichever way, the solutions of such are hard to obtain if they exist. Hence, the quest for effective and reliable methods of solution [1-12]. The model to be considered in this work is the generalized telegraph equation (TE) which is a linear partial differential equation (PDE) describing the current and voltage on an electrical transmission line with x and t as distance and time parameter respectively. The generalized form of the telegraph equation is as follows:

$$\begin{cases} \Omega_{tt} + (a+b)\Omega_{t} + ab\Omega = c^{2}\Omega_{xx} + g(x,t), \ \Omega = \Omega(x,t), \\ \Omega(x,0) = f_{1}(x), \\ \Omega_{t}(x,0) = f_{2}(x), \end{cases}$$
(1.1)

where the constants a,b, and c are real numbers, while  $g(x,t), f_1(x)$ , and  $f_2(x)$  are known functions. The unknown function,  $\Omega = \Omega(x,t)$  to be determined denotes voltage or current through the wire at position, X with respect to time, t. The derivation of (1.1) is contained in [13].

Recently, a good number of solution analysts have deliberated on various techniques for the exact and/or approximate solution to (1.1) [14-24]. This work proposes He's polynomial method for the solution of (1.1) [8-12, 25, 26]. Related researches on communication system, circuit, wave

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

**IOP** Publishing

IOP Conf. Series: Materials Science and Engineering 640 (2019) 012117 doi:10.1088/1757-899X/640/1/012117

transmission, networking and so on include those of [27-28]. It is worth noting that the method being introduced is entwined in terms of applications with numerical analysis, computational finance, stochastic or random differential equations, and so on [29-34].

#### 2. Remark on the HPSM

Let  $\theta$  be an operator (integral or differential), such that:

$$\theta(\nu) = 0. \tag{2.1}$$

Suppose we defined a convex homotopy function, H(c, f) by:

$$H(c,f) = f\theta(c) + (1-f)G(c), \qquad (2.2)$$

such that G(c) is referred to as a functional operator. Hence, we get:

$$H(c,0) = G(c) \text{ and } H(c,1) = \theta(c),$$
 (2.3)

if H(c, f) = 0 is satisfied and a given embedded parameter,  $f \in (0,1]$  is considered. In HPM, f = p is applied as an expanding parameter term to obtain:

$$\begin{cases} c = \nu = \lim_{p \to 1} \left( \sum_{j=0}^{\infty} p^{j} \nu_{j} \right), \\ N(\nu) = \sum_{j=0}^{\infty} p^{j} H_{j}. \end{cases}$$
(2.4)

The approach takes the nonlinear term to be  $N(\nu)$  whenever (2.1) is decomposed, such that  $H_k$ 's are the so-called He's polynomials defined as:

$$H_k\left(\nu_0, \nu_1, \nu_2, \cdots, \nu_k\right) = \frac{1}{k!} \frac{\partial^k}{\partial p^k} \left( N\left(\sum_{j=0}^k p^j \nu_j\right) \right)_{p=0}, \ k \ge 0.$$

$$(2.5)$$

#### 3. The Generalized Telegraph Model and the HPSM (He's Polynomials Solution Method)

The HPSM is applied to the Telegraph equation in (1.1) as follows. Let us re-write (1.1) in integral form, while the two-fold integral operator,  $I_0^t(\cdot)$  is applied accordingly. Thus:

$$\begin{cases} \Omega = \Omega(x,0) + \Omega_t(x,0)t + I_0^t(c^2\Omega_{xx} + g(x,t) - ((a+b)\Omega_t + ab\Omega)), \\ \Omega(x,0) = f_1(x), \Omega_t(x,0) = f_2(x), \ \Omega(x,t) = \Omega. \end{cases}$$
(3.1)

This implies that:

$$\Omega = \underbrace{f_1(x) + f_2(x)t + I_0^t(g(x,t))}_{\text{initial-resultant}} + I_0^t(c^2\Omega_{xx} - ((a+b)\Omega_t + ab\Omega)).$$
(3.2)

In standard HPSM, the series solution is conveyed as:

$$\begin{cases} \Omega = \sum_{i=0}^{\infty} p^{i} \Omega_{i} \\ \Omega = \Omega(x,t), p \to 1. \end{cases}$$
(3.3)

Hence, by homotopy convexity [9, 10] in line with (3.3), we have:

**IOP** Publishing

IOP Conf. Series: Materials Science and Engineering 640 (2019) 012117 doi:10.1088/1757-899X/640/1/012117

$$\begin{cases} \sum_{n=0}^{\infty} p^{n} \Omega_{n} = F(x) + I_{0}^{t} \left( c^{2} \sum_{n=0}^{\infty} p^{n+1} (\Omega_{n})_{xx} - \left( (a+b) \sum_{n=0}^{\infty} p^{n+1} (\Omega_{n})_{t} + ab \sum_{n=0}^{\infty} p^{n+1} (\Omega_{n}) \right) \right), \\ F(x,t) = f_{1}(x) + f_{2}(x)t + I_{0}^{t} \left( g(x,t) \right). \end{cases}$$
(3.4)

Thus, comparing the exponents (powers) of the p's in (3.4), we have:

$$\begin{cases} \Omega_{0} = F(x,t) \\ \Omega_{1} = I_{0}^{t} \left\{ c^{2} \left( \Omega_{0} \right)_{xx} - \left( (a+b) \left( \Omega_{0} \right)_{t} + ab \Omega_{0} \right) \right\} \\ \Omega_{2} = I_{0}^{t} \left\{ c^{2} \left( \Omega_{1} \right)_{xx} - \left( (a+b) \left( \Omega_{1} \right)_{t} + ab \Omega_{1} \right) \right\} \\ \left\{ \Omega_{3} = I_{0}^{t} \left\{ c^{2} \left( \Omega_{2} \right)_{xx} - \left( (a+b) \left( \Omega_{2} \right)_{t} + ab \left( \Omega_{2} \right) \right) \right\} \\ \Omega_{4} = I_{0}^{t} \left\{ c^{2} \left( \Omega_{3} \right)_{xx} - \left( (a+b) \left( \Omega_{3} \right)_{t} + ab \left( \Omega_{3} \right) \right) \right\} \\ \vdots \end{cases}$$

$$(3.5)$$

$$(3.5)$$

$$(3.6)$$

$$\Omega_{j+1} = I_0^t \left\{ c^2 \left( \Omega_j \right)_{xx} - \left( (a+b) \left( \Omega_j \right)_t + ab \left( \Omega_j \right) \right) \right\}, \ j \ge 0.$$

So, the required solution is:

$$\Omega(x,t) = \lim_{p \to 1} \left( \sum_{n=0}^{\infty} p^n \Omega_n \right).$$
(3.7)

### 4. Applications

Let the following linear telegraph equation be considered [15, 24]:

$$\begin{cases} \Omega_{tt} + \Omega_{t} + \Omega = \Omega_{xx}, \ \Omega = \Omega(x, t), \\ \Omega(x, 0) = e^{x}, \\ \Omega_{t}(x, 0) = -e^{x}, \end{cases}$$
(4.1)

with an exact solution of the form:

$$\Omega(x,t) = e^{x-t} \,. \tag{4.2}$$

If (4.1) is compared with (1.1), then we have:

$$(a+b) = ab = c = 1, g(x,t) = 0, f_1(x) = e^x, \text{ and } f_2(x) = -e^x.$$
 (4.3)

So, the recursive relation based on (3.5) and (3.6) is:

$$\begin{cases} \Omega_0 = (1-t)e^x, \\ \Omega_{j+1} = I_0^t \left\{ \left(\Omega_j\right)_{xx} - \left(\left(\Omega_j\right)_t + \left(\Omega_j\right)\right) \right\}, j \ge 0. \end{cases}$$

$$(4.4)$$

Hence,

$$\Omega_0 = (1-t)e^x,$$
  

$$\Omega_1 = I_0^t \left\{ (\Omega_0)_{xx} - ((\Omega_0)_t + (\Omega_0)) \right\} = \frac{t^2 e^x}{2!},$$

$$\Omega_{2} = I_{0}^{t} \left\{ (\Omega_{1})_{xx} - ((\Omega_{1})_{t} + (\Omega_{1})) \right\} = -\frac{t^{3}e^{x}}{3!},$$
  

$$\Omega_{3} = I_{0}^{t} \left\{ (\Omega_{2})_{xx} - ((\Omega_{2})_{t} + (\Omega_{2})) \right\} = \frac{t^{4}e^{x}}{4!},$$
  

$$\Omega_{4} = I_{0}^{t} \left\{ (\Omega_{3})_{xx} - ((\Omega_{3})_{t} + (\Omega_{3})) \right\} = -\frac{t^{5}e^{x}}{5!},$$
  

$$\vdots$$

Thus,

$$\Omega(x,t) = (1-t)e^{x} + \frac{t^{2}e^{x}}{2!} - \frac{t^{3}e^{x}}{3!} + \frac{t^{4}e^{x}}{4!} - \frac{t^{5}e^{x}}{5!} + \cdots$$

$$= \left(1 - t + \frac{t^{2}}{2!} - \frac{t^{3}}{3!} + \frac{t^{4}}{4!} - \frac{t^{5}}{5!} + \cdots\right)e^{x}$$

$$= e^{x-t}.$$
(4.5)

The solution in (4.5) corresponds to those obtained in [14, 19]. Though, the approach contained herein appears simpler and straight forward. The approximate and the exact solutions are graphically displayed in Figure 1 and Figure 2 respectively.



Figure 1: HPSM 6-term Approximate solution



Figure 2: HPSM Exact solution

#### 5. Conclusions

This work has successfully presented the application of the proposed solution method referred to as HPSM to the generalized telegraph equation in terms of approximate-analytical solutions. Closed form solutions of the solved problems were realized with ease, even with less computational time. Though, it may require coupling with other methods for highly nonlinear models. Hence, the HPSM is recommended for nonhomogeneous version of the generalized telegraph equation, and other highly nonlinear models.

#### Acknowledgement

CUCRID section of Covenant University is highly appreciated for all forms of support.

#### References

- [1] Wazwaz A M and Mehanna M S 2010 The combined Laplace–Adomian method for handling singular integral equation of heat transfer, *Int J Nonlinear Sci.*, **10** 248-52
- [2] Ablowitz M J Herbst B M and Schober C 1996 Constance on the numerical solution of the sine-Gordon equation. I: Integrable discretizations and homoclinic manifolds, J. Comput Phys 126 299–314
- [3] Edeki S O Akinlabi G O and Adeosun S A On a modified transformation method for exact and approximate solutions of linear Schrödinger equations, *AIP Conference Proceedings* **1705** 020048
- [4] Akinlabi G O and Edeki S O On Approximate and Closed-form Solution Method for Initialvalue Wave-like Models, *International Journal of Pure and Applied Mathematics* 107(2) 449– 456

- [5] Edeki S O and Akinlabi G O 2017 Coupled method for solving time-fractional navier stokes equation, *International Journal of Circuits, Systems and Signal Processing* **12** 27-34
- [6] Edeki S O Akinlabi G O and Adeosun S A 2016 Analytic and Numerical Solutions of Time-Fractional Linear Schrödinger Equation, *Comm Math Appl*, 7(1) 1-10
- [7] Edeki S O and Akinlabi G O 2017 Zhou Method for the Solutions of System of Proportional Delay Differential Equations, *MATEC Web of Conferences* **125** 02001
- [8] Edeki S O Ugbebor O O and Owoloko E A 2016 He's polynomials for analytical solutions of the Black-Scholes pricing model for stock option valuation, *Proceedings of the World Congress on Engineering 2016*, **II**, WCE 2016, June 29 July 1, 2016, London, U.K.
- [9] Ghorbani A and Nadjfi A S 2007 He's homotopy perturbation method for calculating Adomian's polynomials, *Int. J. Nonlin. Sci. Num. Simul.* **8** (2) 229-332
- [10] He J H Homotopy perturbation method: A new nonlinear analytical technique, *Appl. Math. Comput*, **135** 73-79
- [11] Saberi-Nadjafi J and Ghorbani A He's homotopy perturbation method: an effective tool for solving nonlinear integral and integro-differential equations, *Computers & Mathematics with Applications*, **58**, (2009), 1345-1351.
- [12] Singh J Kumar K and Rathore S 2012 Application of Homotopy Perturbation Transform Method for Solving Linear and Nonlinear Klein-Gordon Equations, *Journal of Information and Computing Science* 7 (2) 131-139
- [13] Srivastava V K Awasthi M K Chaurasia R K and Tamsir M 2013 The Telegraph Equation and Its Solution by Reduced Differential Transform Method, *Modelling and Simulation in Engineering* 2013 746351
- [14] M. Sari, A. Gunay, G. Gurarslan, A Solution to the Telegraph Equation by Using DGJ Method, *International Journal of Nonlinear Science*, 17 (1), (2014), 57-66.
- [15] Gao F and Chi C 2007 Unconditionally stable difference schemes for a one-space-dimensional linear hyperbolic equation, *Appl. Math. Comput.* 187 1272-1276
- [16] Dehghan M and Shokri A 2008 A numerical method for solving the hyperbolic telegraph equation. *Numer. Methods Partial Differential Eq.* **24** 1080-1093
- [17] Aloy R Casaban M C Caudillo-Mata L A and Jodar L 2007 Computing the variable coefficient telegraph equation using a discrete eigenfunctions method, *Comput. Math. Appl.* 54 448-458
- [18] Mohebbi A and Dehghan M High order compact solution of the one-space-dimensional linear hyperbolic equation, *Numer. Methods Partial Differential Eq.* **24** 1222-1235
- [19] Biazar J Ebrahimi H and Ayati Z 2009 An approximation to the solution of telegraph equation by variational iteration method, *Numer. Methods Partial Differential Eq.* **25**(2009) 797-801
- [20] Mohanty R K Jain M K and George K 1996 On the use of high order difference methods for the system of one space second order non-linear hyperbolic equations with variable coefficients J. Comp. Appl. Math. 72 421-431.
- [21] El-Azab M S and El-Gamel M 2007 A numerical algorithm for the solution of telegraph equations, *Appl. Math. Comput.* **190** 757-764
- [22] Mohanty R K Jain M K and Arora O An unconditionally stable ADI method for the linear hyperbolic equation in three space dimensions, *Int. J. Comput. Math.* **79** 133-142
- [23] Dehghan M and Lakestani M 2009 The use of Chebyshev cardinal functions for solution of the second-order one- dimensional telegraph equation, *Numer. Methods Partial Differential Eq.* 25 931-938.
- [24] Saadatmandi A and Dehghan M 2009 Numerical solution of hyperbolic telegraph equation using the Chebyshev Tau method, *Numer*. *Methods Partial Differential Eq* (in press)
- [25] Edeki S O Akinlabi G O and Adeosun M E 2016 Analytical solutions of the Navier-Stokes model by He's polynomials, Lecture Notes in Engineering and Computer Science 2223 16-19
- [26] Edeki S O Akinlabi G O and Adeosun S A 2017 Approximate-analytical solutions of the generalized newell-whitehead-segel model by He's polynomials method, *Lecture Notes in Engineering and Computer Science* 2229 57-59

- [27] Jonathan O Olajide F and Ayo C 2016 Network forensics tools in a mixed-network environment and the adoption of e-voting system in developing countries, *International Journal of Pharmacy* and Technology 8(4) 23115-23118
- [28] Usikalu M R Ilesanmi O R 2018 Network elements of a telecommunication service provider in Nigeria, 2018 IEEE Conference on Technologies for Sustainability, SusTech 8671377
- [29] Edeki S O Ugbebor O O and Owoloko E A 2018 On a dividend-paying stock options pricing model (sopm) using constant elasticity of variance stochastic dynamics, *International Journal* of Pure and Applied Mathematics 106 (4) 1029-1036
- [30] Akinlabi G O and Edeki S O 2017 the solution of initial-value wave-like models via perturbation iteration transform method, *Lecture Notes in Engineering and Computer Science* **2228** 1015-1018.
- [31] Edeki S O Owoloko E A and Ugbebor O O 2015 The modified Black-Scholes model via constant elasticity of variance for stock options valuation, 2015 Progress in Applied Mathematics in Science and Engineering (PIAMSE), AIP Conference Proceedings 1705 4940289
- [32] El-Tawil M A and Tolba A H On the application of mean square calculus for solving random differential equations, *Electronic Journal of Mathematical Analysis and Application* 1 (2) (2013) 202-211
- [33] Edeki S O Ugbebor O O and Owoloko E A 2017 analytical solution of the time-fractional order Black-Scholes model for stock option valuation on no dividend yield basis, *IAENG International Journal of Applied Mathematics* 47 (4) 407-416
- [34] Fakharzadeh A Hesamaeddini E Soleimanivareki M 2015 Multi-step stochastic differential transformation method for solving some class of random differential equations, *Applied Mathematics in Engineering, Management and Technology* **3**(3) 115-123