



Identification of the woody biomasses in Southwest, Nigeria as potential energy feedstocks in thermal power plants for air pollution control

F. B. Elehinafe , O. B. Okedere , J. A. Sonibare & A. O. Mamudu |

To cite this article: F. B. Elehinafe , O. B. Okedere , J. A. Sonibare & A. O. Mamudu | (2021) Identification of the woody biomasses in Southwest, Nigeria as potential energy feedstocks in thermal power plants for air pollution control, Cogent Engineering, 8:1, 1868146, DOI: [10.1080/23311916.2020.1868146](https://doi.org/10.1080/23311916.2020.1868146)

To link to this article: <https://doi.org/10.1080/23311916.2020.1868146>



© 2021 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.



Published online: 08 Jan 2021.



Submit your article to this journal [↗](#)



Article views: 137



View related articles [↗](#)



View Crossmark data [↗](#)



Received: 20 September 2020
Accepted: 18 December 2020

*Corresponding author: F. B. Elehinafe, Department of Chemical Engineering, Covenant University, Ota, Ogun State, Nigeria
E-mail: francis.elehinafe@covenantuniversity.edu.ng

Reviewing editor:
Harvey Arellano-Garcia, Arellano-Garcia, Osun State University, Nigeria

Additional information is available at the end of the article

CHEMICAL ENGINEERING | RESEARCH ARTICLE

Identification of the woody biomasses in Southwest, Nigeria as potential energy feedstocks in thermal power plants for air pollution control

F. B. Elehinafe^{1*}, O. B. Okedere², J. A. Sonibare³ and A. O. Mamudu¹

Abstract: This study identified one hundred samples of woody biomasses in the southwest, Nigerian. The woody biomass samples identified were collected from different saw mills, farms and kitchens in Osun State, Nigeria. The identification of the woody biomass samples was done using literature materials and at a herbarium in the Botany Department, Obafemi Awolowo University, Nigeria, using the vegetative parts of their trees. The identification was carried out to know the numerical availability, the family or English names, the common names and the botanical names of the southwest Nigeria woody biomasses. From the findings, 100 samples of the woody biomasses were identified of 39 families and classes of hard woody biomasses and soft woody biomasses out of which only two are soft and rest are hard. It was discovered that Southwest, Nigeria has the capacity to embark on the massive production of these woody biomasses identified due to her land mass of 77,818 km² comprising 85 forests. In conclusion, serious investment in the propagation of these energy crops would lead to the production of woody masses to serve as feedstocks in thermal power plants for sustainability in power supply, air pollution control and employment opportunities.

Subjects: Power & Energy; Renewable Energy; Clean Tech; Renewable Energy

ABOUT THE AUTHORS

Francis B. Elehinafe holds a PhD degree in Chemical Engineering in Obafemi Awolowo University, Ile-Ife, Osun State, Nigeria. He is currently a lecturer and researcher at Covenant University, Ota, Ogun State, Nigeria. His researcher areas include environmental pollution (monitoring, analysis, modelling and control); waste management; and renewable energy.

Oyetunji B. Okedere's research interest include air pollution (monitoring, analysis, modelling and control), and renewable energy.

Angela O. Mamudu's is a researcher with interest in process optimization; health, safety and environment; plant design; petroleum refining; and petrochemicals.

Ademola J. Sonibare's research interest include air pollution (monitoring, analysis, modelling and control)

PUBLIC INTEREST STATEMENT

Woody biomasses when utilized as fuel discharge carbon recently removed from the air, thereby, hypothetically not expanding GHG levels in the atmosphere. The GHG discharges during the burning of woody biomass are thought to be equivalent to that taken from the atmosphere through the process of photosynthesis during woody biomasses were growing. Woody biomasses as sources of energy do not fluctuate like other renewable sources of energy. For example, solar energy is not available during cloud cover and rainfall, hydro-electricity drops during the dry season. Seasons do not affect the supply of fuel-woods once trees have grown to the peak in sizes, numbers and as the ripe ones are being felled, they are replanted. So, power plants fired using woody feedstocks would never have a down time in operations. Hence, air emissions associated with the combustion of fossil fuels are controlled.

Keywords: Woody biomass; energy feedstocks; thermal conversion; power plants; air pollution control

1. Introduction

Air Pollution is a huge concern faced by the world today and impacts all human beings in so many different ways. The decisions concerning the fight against air pollution should be guided by the understanding that economic development, social development and environmental protection are interdependent and mutually reinforcing components of sustainable development (Ogbonnaya et al., 2007; Ganesan et al., 2015). The growing interest in moving energy generation away from petroleum fuels and toward alternative means is pulling together considerations for the utilization of renewable energy sources to help meet the global energy needs and sustainability of the environment. Thus, the previous decade has witnessed an increasing attention for understanding the roles the renewable energy sources can play in the energy adequacy and a quantum rise in research support into conversion technologies and obtainability as reported by a number of authors (Ganesan et al., 2013; Hoseinzadeh et al., 2020; Javadi et al., 2020; Mason et al., 2009; Siamak et al., 2020).

Nigeria is enriched with adequate renewable sources of energy assets to meet its present and future growth and development prerequisites. Nonetheless, hydropower is the main reasonable resource at present exploited and connected to the network of electricity grid (Nnaji et al., 2010). Sourcing for renewable and sustainable energy development and awareness creation in Nigeria is driven by, in addition to other things, the ongoing increment in fossil fuel prices, the inaccessibility of power supply to the greater part of the Nigerians and the significant losses associated with cost, energy during grid extension and air pollution control. The Nigerian government has put forth attempts through a lot of endeavours, policies and programmes to encourage private investments to go into renewable energy exploitation (Ogbonnaya et al., 2007). Notwithstanding, there are impediments, primarily because of technical and money-related hindrances, that should be addressed for this to be a reality. More importantly, the carbon, sulphur, nitrogen and ash contents of the woody biomasses are insignificant and mild to constitute air pollution (Boluwaji et al., 2019) and the energy contents of woody biomasses are at par with those of the fossils.

In Nigeria, the absolute value of both the woody and non-woody products of forest just as their ecological functions is tremendous however not totally quantifiable (Daniel & Babatunde Hope, 2020). Generally, the forest is underrated in its embedded value within the government economic reckoning of Nigerian woody vegetation assets comprise the thick forest, bushland, woodland, trees on farmlands and plantation. Every one of these different resources adds to production, conservation and protection functions of the national vegetation. In the nation, studies have indicated that forest reserves extend to 10 million hectares, which represents about 10% of a land region of roughly 96.2 million hectares (Adelakun & Olugbade, 2019). Nigeria is well geographically located for maximum rainfall and temperature throughout the year for plant growth thus for woody biomass production (Agboola, 1979). There are a lot of woody biomasses that could be used as feedstocks for power generation in this part of the globe.

Southwestern Nigeria has eighty-five forest reserves with a total area cover of 842,499 hectares (Adelakun & Olugbade, 2019). The climate of the region is tropical naturally and it is portrayed by dry and wet seasons. The temperature ranges somewhere in the range of 21°C and 34°C while the annual precipitation ranges from 1500 mm to 3000 mm (Adelakun & Olugbade, 2019). The wet season is related with the southwest monsoon winds from the Atlantic Ocean while the dry season is associated with the northeast trade wind from the Sahara Desert. Agboola (1979) reported that the vegetation in Southwest, Nigeria comprises mangrove forest and fresh water swamp at the belt region, the low land in primary forest zone stretching inland to Ogun State and part of Ondo State while the secondary forest is towards the northern borderline where southern savannah and derived savannah likewise exist. This forest area of the southwest is enough to meet the energy

need of Nigeria through woody biomass production for power generation if serious attention is paid to it by the government.

Consumption of petroleum fuels and other fossils releases carbon put away underneath the crust of the earth for centuries into the air, a crucial part of the environment as ozone-depleting substances called greenhouse gases (GHG). Be that as it may, these gasses are answerable for a global temperature alteration (Demirbas, 2005; Odunlami Olayemi et al., 2018; Ganesan et al., 2015). Then again, woody biomass when utilized as fuel discharge carbon recently removed from the air, thereby, hypothetically not expanding GHG levels in the atmosphere. The GHG discharges during the burning of woody biomass are thought to be equivalent to that taken from the atmosphere through the process of photosynthesis during woody biomasses were growing (Boluwaji et al., 2019; Demirbas, 2005), particularly on account of agricultural residues and forest (Biagini et al., 2006). In any case, a general life cycle investigation has demonstrated a rise in GHG emissions for woody biomass usage for biofuels (ICF, 2009). The rise in GHG emissions was the consequence of the utilization of fossils for energy production during different tasks engaged with, in supply-chain logistics (Nitschke & Innes, 2013). Then again, revealed that woody biomass gathering from sustainably managed forests really diminishes net GHG emissions, in this way, making it carbon negative.

Fossil-fired thermal plants, unlike biomass-fired thermal plants, produce large quantities of Sulphur IV oxide (SO₂) and nitrogen oxides (NO_x), the basic deleterious emissions in the acid rain. Acid rain increases the pH of water bodies, and damages the vegetation and seaside biological systems (Elehinafe Francis et al., 2017; Piotr, 2020; Trang et al., 2020). SO₂ and NO_x engender particulate formation. It is reported by US EPA in 2004 that ozone (smog) and nitrates are formed with NO_x as precursors. Ozone mars the functions of lungs and diminishes the yields of numerous economically significant agricultural plants. Nitrates in precipitation make water bodies over-enriched, causing algal sprouts that kill aquatic and hamper biodiversity.

Woody biomasses as sources of energy do not fluctuate like other renewable sources of energy. For example, solar energy is not available during cloud cover and rainfall, hydro-electricity drops during the dry season. Seasons do not affect the supply of fuelwoods once trees have grown to the peak in sizes, numbers and as the ripe ones are being felled, they are replanted (Balat et al., 2009). So, power plants fired using woody feedstocks would never have a down time in operations. Hence, air emissions associated with the combustion of fossil fuels are controlled. Considering woody biomass potentials Nigeria has and the environment, there is need to identify, as many as possible, the available woody biomasses in terms of trees in Nigeria to the knowledge of investors, government and researchers.

2. Methodology

2.1. The study area

The study area is Southwest Nigeria which comprises Oyo, Ekiti, Ondo, Osun, Ogun, and Lagos States (Figure 1). The area is between Latitudes 6° 21¹ and 8° 37¹ N and Longitudes 2° 31¹ and 6° 00¹ East. The study area has an approximate land area of about 77,818 km² and 85 forest reserves with a forest area covering 842,499 hectares as reported by Faleyimu and Agbeja (2012). The climate of Southwest Nigeria is tropical and it is characterized by dry and wet seasons. The temperature ranges between 21°C and 34°C while the annual rainfall ranges between 1500 mm and 3000 mm (Agboola, 1979). The wet season is connected with the Southwest monsoon winds from the Atlantic Ocean and the dry season is associated with the northeast trade winds from the sahara desert. According to the report of Oluseyi (2014), the vegetation in Southwest, Nigeria is made up of fresh water swamp and mangrove forest at the belts, the low land in forest stretches inland to Ogun and part of Ondo state while the secondary forest is towards the northern boundary where derived and southern Savannah exist.

Figure 1. Vegetation map of South-West Nigeria (Izuchukwu et al., 2020).



2.2. Procedure for identification woody biomasses in Southwest, Nigeria

The woody biomass samples identified in this work were collected from different saw mills, farms and kitchens in Osun State, Nigeria. One hundred (100) samples of different common woody biomasses were gotten. Literature materials (Abdulrahman et al., 2006; Erakhrumen et al., 2010; Oladunmoye & Kehinde, 2011; Wahab et al., 2013) were used to identify many of the woody biomasses in translating the vernacular name to the common, family or English and botanical names as well as the woody types—hard or soft. Flowers and fruits as well as vegetative parts of the rest whose vernacular names could not be traced to the common and botanical names in the literature, were taken to the herbarium of Botany Department, Obafemi Awolowo University, Ile-Ife for identification. The woody biomasses were classified into hard woody biomasses and soft woody biomasses.

3. Results and discussion

The results of the identification, nomenclature and classification are presented in Table 1. The identification was carried out to know the common names, the family or English names and the botanical names of the southwest Nigeria woody biomasses. The same categories of woody biomasses are processed at all the saw-mills visited. This is corroborated by the work of Olufemi et al. (2012) that the same kinds of woody biomasses are present in southwest Nigeria. One hundred samples of the woody biomasses identified were grouped into 39 families and classified into hard woody biomasses and soft woody biomasses out of which only two are soft and rest are hard.

From the numerical distribution of the woody biomasses based on families, the woody biomasses comprise 1 of *Bombaceae*, *Rhizophoraceae*, *Poaceae*, *Myrtaceae*, *Annonaceae*, *Gentianaceae*, *Myristicaceae*, *Boraginaceae*, *Lecythidaceae*, *Phyllanthaceae*, *menispermaceae*, *Vitaceae*, *Urticaceae*, *Pinaceae*, *Asteraceae*, *Rosaceae*, *Fagaceae*, *Olacaceae* and *Acanthaceae*; 2 of *Clusiaceae*, *Rubiaceae*, *Lamiaceae*, *Irvingiaceae*, *Sapotaceae*, *leguminoceae*, *Bignoniaceae* and *Sapindaceae*; 3 of *Sterculiaceae*, *Apocynaceae*; 4 of *Arecaceae* and *Combretaceae*; 5 of *Meliaceae* and *Euphorbiaceae*; 6 of *Moraceae*; 7 of *Rutaceae*; 11 of *Malvaceae* and 14 of *Fabaceae*. Ninety-eight of them are hard woody biomasses while two are soft woody biomasses (*Bombax buonopozense* from the family of *Malvaceae* and *Pinus ponderosa* from the family of *Pinaceae*). From the identification and nomenclature of the woody biomasses, the *Fabaceae* family is most abundant, 14%, followed by the *Malvaceae* family, 11%. According to Richardson et al. (2011), they have many species, grow and mature faster than other families. They are also resistant to climatic and edaphic factors. *Fabaceae* and *Malvaceae* families are known for their naturalization and invasion potential for they are mostly

Table 1. Woody biomasses identified in Southwest, Nigeria

| S/N | Botanical name | Family | Common name | Wood type |
|-----|----------------------------------|----------------|----------------|--------------------|
| 1 | <i>Adansonia digitata</i> | Malvaceae | Baobab | Hard woody biomass |
| 2 | <i>Afrormosia elata</i> | Fabaceae | Shedum | Hard woody biomass |
| 3 | <i>Albizia gummifera</i> | Fabaceae | Ayinre Banaban | Hard woody biomass |
| 4 | <i>Anacardium occidentale</i> | Anacardiaceae | Cashew | Hard woody biomass |
| 5 | <i>Anogeissus leiocarpus</i> | Combretaceae | Orin Dudu | Hard woody biomass |
| 6 | <i>Antiaris africana</i> | Moraceae | Oriro | Hard woody biomass |
| 7 | <i>Anthocleista vogelii</i> | Gentianaceae | Sapo sapo | Hard woody biomass |
| 8 | <i>Antrocaryon micraster</i> | Anacardiaceae | Akikogbon | Hard woody biomass |
| 9 | <i>Artocarpus altilis</i> | Moraceae | Bread fruit | Hard woody biomass |
| 10 | <i>Asteromyrtus symphyocarpa</i> | Myrtaceae | Waria | Hard woody biomass |
| 11 | <i>Alstonia boonei</i> | Apocynaceae | Ahun funfun | Hard woody biomass |
| 12 | <i>Azadirachta indica</i> | Meliaceae | Dongoyaro | Hard woody biomass |
| 13 | <i>Berlinia grandifolia</i> | Fabaceae | Itako | Hard woody biomass |
| 14 | <i>Blighia sapida</i> | Sapindaceae | Isin | Hard woody biomass |
| 15 | <i>Bombax buonopozense</i> | Malvaceae | Ponpola | Hard woody biomass |
| 16 | <i>Brachystegia leonensis</i> | Fabaceae | Ako | Hard woody biomass |
| 17 | <i>Byraria marginata</i> | Acanthaceae | Eso | Hard woody biomass |
| 18 | <i>Cassia fistula</i> | Fabaceae | Cassia | Hard woody biomass |
| 19 | <i>Ceiba pentandra</i> | Malvaceae | Araba | Hard woody biomass |
| 20 | <i>Celtis zenkeri</i> | Ulmaceae | Ita | Hard woody biomass |
| 21 | <i>Chasmanthera dependens</i> | Menispermaceae | Ato | Hard woody biomass |
| 22 | <i>Chrysophyllum africanum</i> | Sapotaceae | Agbalumo | Hard woody biomass |
| 23 | <i>Cissus adenopoda</i> | Vitaceae | Gbolagbola | Hard woody biomass |
| 24 | <i>Citrus limon</i> | Rutaceae | Lemon | Hard woody biomass |
| 25 | <i>Citrus sinensis</i> | Rutaceae | Osan pupa | Hard woody biomass |
| 26 | <i>Citrus paradisi</i> | Rutaceae | Osan wewe | Hard woody biomass |

(Continued)

Table 1. (Continued)

| S/N | Botanjcal name | Family | Comman name | Wood type |
|-----|-----------------------------------|---------------|--------------------------|--------------------|
| 27 | <i>Citrus aurantifolia</i> | Rutaceae | Grape | Hard woody biomass |
| 28 | <i>Citrus medica</i> | Rutaceae | Tanjarini | Hard woody biomass |
| 29 | <i>Cleistopholis patens</i> | Annonaceae | Apako | Hard woody biomass |
| 30 | <i>Cocos nucifera</i> | Areaceae | Agbon | Hard woody biomass |
| 31 | <i>Cola millenii</i> | | Sterculiaceae | Hard woody biomass |
| 32 | <i>Cola acuminata</i> | | Sterculiaceae | Hard woody biomass |
| 33 | <i>Cola nitida</i> | | Sterculiaceae Obi Gbanja | Hard woody biomass |
| 34 | <i>Cordia milleni</i> | Boraginaceae | Omoh | Hard woody biomass |
| 35 | <i>Crassocephalum biafrae</i> | Asteraceae | Gbolagi | Hard woody biomass |
| 36 | <i>Cylicodiscus gabunensis</i> | Fabaceae | Okan | Hard woody biomass |
| 37 | <i>Daniella oliveri</i> | Leguminaceae | Iyaa | Hard woody biomass |
| 38 | <i>Daniella ogea</i> | Leguminaceae | Asunwole | Hard woody biomass |
| 39 | <i>Delonix regia</i> | Fabaceae | Panseke | Hard woody biomass |
| 40 | <i>Diospyros crassiflora</i> | Fabaceae | Ebony | Hard woody biomass |
| 41 | <i>Elaeis guineensis</i> | Areaceae | Ope Igbo | Hard woody biomass |
| 42 | <i>Entanda gigas</i> | Fabaceae | Agbaa | Hard woody biomass |
| 43 | <i>Entadrophragma cylindricum</i> | Meliaceae | Sapelle Mahogany | Hard woody biomass |
| 44 | <i>Eucalyptus marginata.</i> | Myrtaceae | Jarrah | Hard woody biomass |
| 45 | <i>Ficus thionningii</i> | Moraceae | Odanko | Hard woody biomass |
| 46 | <i>Ficus carica</i> | Moraceae | Opoto | Hard woody biomass |
| 47 | <i>Ficus mucuso</i> | Moraceae | Obobo | Hard woody biomass |
| 48 | <i>Funtumia elastica</i> | Apocynaceae | Ire | Hard woody biomass |
| 49 | <i>Garcinia kola</i> | Clusaceae | Orogbo | Hard woody biomass |
| 50 | <i>Glirocidia sepium</i> | Fabaceae | Agunmoniye | Hard woody biomass |
| 51 | <i>Gmelina arborea</i> | Lamiaceae | Milaina | Hard woody biomass |
| 52 | <i>Hevea brasiliensis</i> | Euphorbiaceae | Para Rubber | Hard woody biomass |

(Continued)

| S/N | Botanical name | Family | Common name | Wood type |
|-----|----------------------------------|----------------|------------------|--------------------|
| 53 | <i>Hildegardia barteri</i> | Malvaceae | Okurugbedu | Hard woody biomass |
| 54 | <i>Hymenocardia acida</i> | Phyllanthaceae | Kampalaga | Hard woody biomass |
| 55 | <i>Irvingia grandifolia</i> | Irvingiaceae | Karakoro | Hard woody biomass |
| 56 | <i>Irvingia excelsa</i> | Irvingiaceae | Oro (wild Mango) | Hard woody biomass |
| 57 | <i>Isoberlina doka</i> | Fabaceae | Babo | Hard woody biomass |
| 58 | <i>Khaya ivorensis</i> | Meliaceae | Ogano | Hard woody biomass |
| 59 | <i>Lecaniodiscus cupanioides</i> | Sapindaceae | Isin | Hard woody biomass |
| 60 | <i>Lophira lanceolata</i> | Ochnaceae | Pahan | Hard woody biomass |
| 61 | <i>Lovoa trichlioides</i> | Meliaceae | Tiger wood | Hard woody biomass |
| 62 | <i>Macaranga barteri</i> | Euphorbiaceae | Asasa | Hard woody biomass |
| 63 | <i>Mangifera indica</i> | Anacardiaceae | Mango | Hard woody biomass |
| 64 | <i>Mansonia altissima</i> | Malvaceae | Mansonia | Hard woody biomass |
| 65 | <i>Milicia excelsa</i> | Moraceae | Iroko | Hard woody biomass |
| 66 | <i>Mitragyna ciliata</i> | Rubiaceae | Abora | Hard woody biomass |
| 67 | <i>Musanga cecropioides</i> | Urticaceae | Alagbao | Soft woody biomass |
| 68 | <i>Napoleona vogelii</i> | Lecythidaceae | Ito | Hard woody biomass |
| 69 | <i>Nauclea diderrichii</i> | Rubiaceae | Opepe | Hard woody biomass |
| 70 | <i>Nesogordonia paparivera</i> | Malvaceae | Oro | Hard woody biomass |
| 71 | <i>Newbouldia laevis</i> | Bignoniaceae | Akoko | Hard woody biomass |
| 72 | <i>Parkia biglobosa</i> | Fabaceae | Irugba | Hard woody biomass |
| 73 | <i>Percuguaria daemia</i> | Apocynaceae | Koleagbe | Hard woody biomass |
| 74 | <i>Phoenix dactylifera</i> | Arecaceae | Date Palm | Hard woody biomass |
| 75 | <i>Pinus ponderosa</i> | Pinaceae | Pine | Hard woody biomass |
| 76 | <i>Piptadeniasrum africanum</i> | Rosaceae | Agboin | Hard woody biomass |
| 77 | <i>Poga oleosa</i> | Rhizophoraceae | Ekú- Ijebu | Hard woody biomass |
| 78 | <i>Prunus dulcis</i> | Rutaceae | Fruit | Hard woody biomass |

(Continued)

Table 1. (Continued)

| S/N | Botanical name | Family | Common name | Wood type |
|-----|---------------------------------|----------------|-------------|--------------------|
| 79 | <i>Pterygota macrocarpa</i> | Malvaceae | Poroporo | Hard woody biomass |
| 80 | <i>Pterocarpus erinaceus</i> | Fabaceae | Gbigbigbi | Hard woody biomass |
| 81 | <i>Pterocarpus osun</i> | Leguminoceae | Osun Pupa | Hard woody biomass |
| 82 | <i>Pycnanthus angolensis</i> | Myristicaceae | Akomu | Hard woody biomass |
| 83 | <i>Quecus robur</i> | Fagaceae | Ayan Iroko | Hard woody biomass |
| 84 | <i>Raphia africana</i> | Arecaceae | Ayan Oguro | Hard woody biomass |
| 85 | <i>Ricinodendron heudelotti</i> | Eurphorbiaceae | Orunmodo | Hard woody biomass |
| 86 | <i>Spondias mombin</i> | Anacardiaceae | Iyeye | Hard woody biomass |
| 87 | <i>Sterculia rhinopetala</i> | Malvaceae | Aye | Hard woody biomass |
| 88 | <i>Strombosia pustulata</i> | Olacaceae | Itako | Hard woody biomass |
| 89 | <i>Strychnos spinosa</i> | Loganiaceae | Atako | Hard woody biomass |
| 90 | <i>Swietenia macrophylla</i> | Meliaceae | Mahogany | Hard woody biomass |
| 91 | <i>Symphonia globulifera</i> | Clusaceae | Okilolo | Hard woody biomass |
| 92 | <i>Tectona grandis</i> | Lamiaceae | Teak | Hard woody biomass |
| 93 | <i>Terminalia glaucescens</i> | Combretaceae | Idi | Hard woody biomass |
| 94 | <i>Terminalia ivorensis</i> | Combretaceae | Idigbo | Hard woody biomass |
| 95 | <i>Terminalia superba</i> | Combretaceae | Afara | Hard woody biomass |
| 96 | <i>Theobroma cacao</i> | Malvaceae | Cocoa | Hard woody biomass |
| 97 | <i>Triplochiton scleroxylon</i> | Malvaceae | Obeche | Hard woody biomass |
| 98 | <i>Uapaca heudelotii</i> | Euphorbiaceae | Akun | Hard woody biomass |
| 99 | <i>Vitellaria paradoxa</i> | Sapotaceae | Emi | Hard woody biomass |
| 100 | <i>Zanthoxylum leprieurii</i> | Rutaceae | Ata | Hard woody biomass |

leguminous plants (Richardson et al., 2011). They have nodules in their roots that house nitrogen-fixing bacteria. So, the soil on which they grow is always fertile. By these, in the absence of deforestation, southwest, Nigeria would never run out of supply of feedstocks to power plants if their utilization as energy carriers, is given attention by the governments and private investors.

Most of the woody biomasses identified are hard, 98%, for they are much more abundant than soft ones in the region. Woody biomasses contain varying amounts of cellulose, hemicellulose, lignin, proteins, simple sugars, starches and small amounts of lipids. They additionally contain

inorganic components and a small amount of water. Among these constituents, hemicellulose, lignin, and cellulose are the three significant constituents with respect to energy or heating value of any woody biomass (Zhang et al., 2010). Lignocellulose is the blend of hemicelluloses, lignin, and cellulose, which involves around half of the plant matter manufactured during photosynthesis and make up the most bountiful renewable natural asset on earth. Cellulose, lignin and hemicelluloses are sturdily interlinked in lignocelluloses and are bonded chemically, by covalent cross-linkages. The biggest part of any lingo-cellulosic material is cellulose, trailed by hemicellulose and then lignin (Zhang et al., 2010). Though hemicellulose and cellulose are macromolecules built from various sugars; lignin is a polymer aromatic in nature produced from phenylpropanoid forerunners.

There are significant differences between hard woody biomasses and soft woody biomasses according to their compositions. Hard woody biomasses are on average about 10% denser than softwoods due to higher lignin and less moisture contents (Braaten & Sellers, 2013). These make hardwoods more useful for energy generation. Southwest, Nigeria is highly favoured naturally with hard woody biomasses to run her thermal plants on renewable feedstocks for air pollution control.

4. Conclusion

Considering the harmful effects of fossil fuels; the advantages of renewable energy sources and the challenges the world faces to have sustainable energy production and environment while reviewing the literature, and the results obtained in this study, i.e., the numerous woody biomasses of families of *Bombaceae*, *Rhizophoraceae*, *Poaceae*, *Myrtaceae*, *Annonaceae*, *Gentianaceae*, *Myristicaceae*, *Boraginaceae*, *Lecythidaceae*, *Phyllanthaceae*, *menispermaceae*, *Vitaceae*, *Urticaceae*, *Pinaceae*, *Asteraceae*, *Rosaceae*, *Fagaceae*, *Olacaceae* and *Acanthaceae*; *Clusiaceae*, *Rubiaceae*, *Lamiaceae*, *Irvingiaceae*, *Sapotaceae*, *leguminoceae*, *Bignoniaceae* and *Sapindaceae*; *Sterculiaceae*, *Apocynaceae*; *Arecaceae* and *Combretaceae*; *Meliaceae* and *Euphorbiaceae*; *Moraceae*; *Rutacaceae*; *Malvaceae* and *Fabaceae* are available in southwest, Nigeria. Hence, the following conclusions were arrived at: the woody biomasses identified are many enough to be termed energy crops in the southwest, Nigeria for them to receive attention from investors and governments; the nation has the land mass that can support massive production of these woody biomasses identified. Employment opportunities would also be created through investment in the propagation of these energy crops in the region. It is expedient that the stakeholders look into these for the attendant benefits.

Acknowledgements

The authors would like to thank Covenant University Centre for Research Innovation and Discovery (CUCRID), Ota, Nigeria for its support in making the publication of this research possible.

Funding

The authors received no direct funding for this research.

Author details

F. B. Elehinafe¹
E-mail: francis.elehinafe@covenantuniversity.edu.ng
O. B. Okedere²
J. A. Sonibare³
A. O. Mamudu¹

ORCID ID: <http://orcid.org/0000-0001-9680-0886>

¹ Department of Chemical Engineering, Covenant University, Ota, Ogun State, Nigeria.

² Faculty of Engineering, Osun State University, Osogbo, Nigeria.

³ Department of Chemical Engineering, Obafemi Awolowo University, Ile-Ife, Nigeria.

Citation information

Cite this article as: Identification of the woody biomasses in Southwest, Nigeria as potential energy feedstocks in thermal power plants for air pollution control, F. B. Elehinafe, O. B. Okedere, J. A. Sonibare & A. O. Mamudu, *Cogent Engineering* (2021), 8: 1868146.

References

- Abdulrahman, A. A., Fajemiroye, O. J., & Oladele, F. A. (2006). Ethnobotanical study of economic trees: Uses of trees as timbers and fuelwoods in ilorin emirate of Kwara State, Nigeria. *Ethnobotanical Leaflets*, 1(1), Article 13. <https://opensiuc.lib.siu.edu/ebf/vol2006/iss1/13>
- Adelakun, G. I., & Olugbade, A. N. (2019). Analysis of pattern and extent of deforestation In Akure Forest Reserve, Ondo State, Nigeria. *Journal of Environmental*.
- Agboola, S. A. (1979). *An agricultural atlas of Nigeria*. Oxford University Press. <https://global.oup.com/academic/product/agricultural-atlas-of-nigeria-9780195754087?lang=ençse>
- Balat, M., Balat, M., Kirtay, E., & Balat, H. (2009). Main routes for the thermo-conversion of biomass into fuels and chemicals. Part 1: Pyrolysis systems. *Energy Conversion and Management*, 50(12), 3147–3157. <https://doi.org/10.1016/j.enconman.2009.08.014>
- Biagini, E., Barontini, F., & Tognotti, L. (2006). Devolatilization of biomass fuels and biomass components studied by TG/FTIR technique. *Industrial & Engineering Chemistry Research*, 45(13), 4486–4493. <https://doi.org/10.1021/ie051404>
- Boluwaji, E. F., Babatunde, O. O., Olayemi, O., Elizabeth, A. O. T., Onose, M. A., & Ademola, S. J. (2019). Comparative study of non-metallic contents of sawdust of different wood species and coal species in

- Nigeria. *Petroleum and Coal*, 61(5), 1183–1189. https://www.vurup.sk/wp-content/uploads/2019/10/PC-X-2019_Elehinofe-99_rev1.pdf
- Braaten, R. W., & Sellers, T. G. (2013). *Prince Edward Island wood chip-fired boiler performance. Division report ERL 92-43 (TR)*. Energy Research Laboratories, Energy, Mines and Resources Canada.
- Daniel, A. A., & Babatunde Hope, O. J. O. (2020). Species Diversity, volume determination and structure of protected forests for in-situ biodiversity conservation. *International Journal of Conservation Science*, 11(1), 133–144. http://ijcs.ro/public/IJCS-20-13_Agbelade.pdf
- Demirbas, A. (2005). Potential applications of renewable energy sources, biomass combustion problems in boiler power systems and combustion related environmental issues. *Progress in Energy and Combustion Science*, 31(2), 171–192. <https://doi.org/10.1016/j.pecs.2005.02.002>
- Elehinafe Francis, B., Okedere Oyetunji, B., Fakinle Bamidele, S., & Sonibare Jacob, A. (2017). Assessment of sawdust of different wood species in Southwestern Nigeria as source of energy. *Energy Sources, Part A: Recovery, Utilization, and Environmental Effects*, 39, 18. <https://doi.org/10.1080/15567036.2017.1384869>.
- Energy Information Administration. 2007b. *Renewables and alternate fuels, wood and wood waste*. Retrieved April 20, 2015, from <http://www.eia.doe.gov/cneaf/solar/renewables/page/wood/wood.html>.
- Erakhrumen, A. A., Ogunsanwo, O. Y., & Ajewole, O. I. (2010). Assessment of some other traditional uses of accepted agroforestry fuelwood species in akinyele and ido local Government Areas, Oyo State, Nigeria.. *International Journal of Social Forestry*, 3(1), 49–65. <http://ir.library.ui.edu.ng/handle/123456789/3409>
- Faleyimu, O. I., & Agbeja, B. O. (2012). Constraints to forest policy implementation in the Southwest Nigeria: Causes, Consequences and Cure. *Resources and Environment*, 2(2), 37–44. <https://doi.org/10.5923/j.re.20120202.06>.
- Ganesan, P., Rajakarunakaran, S., Thirugnanasambandam, M., & Devaraj, D. (2015). The influence of floor tile colour on lighting and energy saving. In C. Kamalakannan, L. Suresh, S. Dash, & B. Panigrahi (Eds.), *Power electronics and renewable energy systems. lecture notes in electrical engineering* (Vol. 326). Springer. https://doi.org/10.1007/978-81-322-2119-7_18.
- Ganesan, P., Rajakarunakaran, S., Thirugnanasambandam, M., & Devaraj, D. (2013). Estimation of energy and carbon saving potential in industrial lighting system. *Advanced Materials Research*, 2013(768), 272–279. <https://doi.org/10.4028/www.scientific.net/AMR.768.272>.
- Hoseinzadeh, S., Roya Yargholi Hamed Kariman, P., & Heyns, S. (2020). Exergo-economic analysis and optimization of reverse osmosis desalination integrated with geothermal energy, 39(5). <https://doi.org/10.1002/ep.13405>.
- ICF. (2009). *Lifecycle greenhouse gas emissions due to increased biofuel production*. Fairfax. Retrieved June 2013, from <http://epa.gov/oms/renewablefuels/rfs2-peer-reviewmodel>
- Izuchukwu, E. N., Chuks, A. P., & Peter, I. K. (2020). Sustainable development through green-themed environmental impact assessment: Lessons for developing countries. *Journal of Environment and Earth Science*, 10(5), 2020.
- Javadi, M. A., Hoseinzadeh, S., Ghasemiasl, R., Heyns, P. S., & Chamkha, A. J. (2020). Sensitivity analysis of combined cycle parameters on exergy, economic, and environmental of a power plant. *Journal of Thermal Analysis and Calorimetry*, 139(1), 519–525. <https://doi.org/10.1007/s10973-019-08399-y>
- Mason, C. L., Lippke, B. R., & Zobrist, K. W. (2009). *Wood to energy in Washington: Imperatives, opportunities, and obstacles to progress*. Report to the Washington State Legislature. https://www.ruraltech.org/pubs/reports/2009/wood_to_energy/pdf/Wood_to_Energy_full_report.pdf
- Nitschke, C. R., & Innes, J. L. (2013). Integrating climate change into forest management in south-central British Columbia: An assessment of landscape vulnerability and development of a climate-smart framework. *Forest Ecology and Management*, 256(3), 313–327. <https://doi.org/10.1016/j.ecolmodel.2007.07.026>
- Nnaji, C. E., Uzoma, C. C., & Chukwu, J. O. (2010). The role of renewable energy resources in poverty alleviation and sustainable development in Nigeria. *Continental Journal Social Sciences*, 3, 31–37. <file:///C:/Users/user/Downloads/ren.Engyrole.pdf>
- Odunlami Olayemi, A., Elehinafe Francis, B., Oladimeji Temitayo, E., Fajobi Muiywa, A., Okedere Oyetunji, B., & Fakinle Bamidele, S. (2018). Implications of lack of maintenance of motorcycles on ambient air quality. *IOP Conf. Series: Materials Science and Engineering*, 413, 012055. <https://doi.org/10.1088/1757-899X/413/1/012055>
- Ogbonnaya, I. O., Chikuni, E., & Govender, P. (2007). *Prospect of wind energy in Nigeria*. Accessed October 2020, from http://active.cput.ac.za/energy/web/duel/papers/2007/0230_Okoro.pdf.
- Oladunmoye, M. K., & Kehinde, F. Y. (2011). Ethno-botanical survey of medicinal plants used in treating viral infections among yoruba tribe of South Western Nigeria. *African Journal of Microbiology Research*, 5(19), 2991–3004. <https://doi.org/10.5897/AJMR10.004>
- Olufemi, B., Akindeni, J. O., & Olaniran, S. O. (2012). Lumber recovery efficiency among selected sawmills in Akure, Nigeria. *Drupa Industriga*, 63(1), 15–18. <https://doi.org/10.5552/drind.2012.1111>
- Oluseyi, A. O. (2014). The potential for wind energy in Nigeria. *Wind Engineering*, 34(3), 303–311. <https://doi.org/10.1260/0309-524X.34.3.303>
- Piotr, B. F. (2020). New technologies and innovative solutions in the development strategies of energy enterprises. *HighTech and Innovation Journal*, 1(2). <https://doi.org/10.28991/HIJ-2020-01-02-01>.
- Richardson, D. M., Carruthers, J., Hui, C., Impson, F. A. C., Miller, J. T., Robertson, M. P., Rouget, M., Le Roux, J. J., & Wilson, J. R. U. (2011). Human-mediated introductions of Australian Acacia species—a global experiment in biogeography. *Diversity and Distributions*, 17(5), 771–787. <https://doi.org/10.1111/j.1472-4642.2011.00824.x>
- Siamak, H., Hadi, G. M., & Stephan, H. (2020). Application of hybrid systems in solution of low power generation at hot seasons for micro hydro systems. *Renewable Energy*, 160, 323–332. <https://doi.org/10.1016/j.renene.2020.06.149>
- Trang, G. T. T., Linh, N. H., Linh, N. T. T., & Kien, P. H. (2020). The study of dynamics heterogeneity in SiO₂ liquid. *HighTech and Innovation Journal*, 1(1), 1–7. <https://doi.org/10.28991/HIJ-2020-01-01-01>
- Wahab, M. K. A., Olalekan, K. K., & Alarape, A. A. (2013). Human activities relating to some economic trees in two local government councils, Osun State, Nigeria. *International Journal of Agric Science*, 3(4), 316–322. file:///C:/Users/user/Downloads/Human_activities_relatng_to_some_econom.pdf
- Zhang, L., Xu, C., & Champagne, P. (2010). Overview of recent advances in thermo-chemical conversion of biomass. *Energy Conversion and Management*, 51(5), 969–982. <https://doi.org/10.1016/j.enconman.2009.11.038>



© 2021 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format.

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.



***Cogent Engineering* (ISSN: 2331-1916) is published by Cogent OA, part of Taylor & Francis Group.**

Publishing with Cogent OA ensures:

- Immediate, universal access to your article on publication
- High visibility and discoverability via the Cogent OA website as well as Taylor & Francis Online
- Download and citation statistics for your article
- Rapid online publication
- Input from, and dialog with, expert editors and editorial boards
- Retention of full copyright of your article
- Guaranteed legacy preservation of your article
- Discounts and waivers for authors in developing regions

Submit your manuscript to a Cogent OA journal at www.CogentOA.com

