

# A Study of Noise Pollution Measurements and Possible Effects on Public Health in Ota Metropolis, Nigeria

Pelumi E. Oguntunde<sup>1\*</sup>, Hilary I. Okagbue<sup>1</sup>, Omoleye A. Oguntunde<sup>2</sup>, Oluwole O. Odetunmibi<sup>1</sup>

<sup>1</sup>Department of Mathematics, Covenant University, Ogun State, Ota, Nigeria; <sup>2</sup>Department of Business Management, Covenant University, Ogun State, Ota, Nigeria

## Abstract

**Citation:** Oguntunde PE, Okagbue HI, Oguntunde OA, Odetunmibi OO. A Study of Noise Pollution Measurements and Possible Effects on Public Health in Ota Metropolis, Nigeria. Open Access Maced J Med Sci. 2019 Apr 30; 7(8):1391-1395. <https://doi.org/10.3889/oamjms.2019.234>

**Keywords:** Environmental toxicity; Distribution; Nigeria; Noise Pollution; Public health

**\*Correspondence:** Pelumi E. Oguntunde, Department of Mathematics, Covenant University, Ogun State, Ota, Nigeria. E-mail: pelumi.oguntunde@covenantuniversity.edu.ng

**Received:** 04-Feb-2019; **Revised:** 21-Mar-2019; **Accepted:** 22-Mar-2019; **Online first:** 29-Apr-2019

**Copyright:** © 2019 Pelumi E. Oguntunde, Hilary I. Okagbue, Omoleye A. Oguntunde, Oluwole O. Odetunmibi. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**Funding:** This research received financial support from the Covenant University

**Competing Interests:** The authors have declared that no competing interests exist

**BACKGROUND:** Noise pollution has become a major environmental problem leading to nuisances and health issues.

**AIM:** This paper aims to study and analyse the noise pollution levels in major areas in Ota metropolis. A probability model which is capable of predicting the noise pollution level is also determined.

**METHODS:** Datasets on the noise pollution level in 41 locations across Ota metropolis were used in this research. The datasets were collected thrice per day; morning, afternoon and evening. Descriptive statistics were performed, and analysis of variance was also conducted using Minitab version 17.0 software. Easy fit software was however used to select the appropriate probability model that would best describe the dataset.

**RESULTS:** The noise levels are way far from the WHO recommendations. Also, there is no significant difference in the effects of the noise pollution level for all the times of the day considered. The log-logistic distribution provides the best fit to the dataset based on the Kolmogorov Smirnov goodness of fit test.

**CONCLUSION:** The fitted probability model can help in the prediction of noise pollution and act as a yardstick in the reduction of noise pollution, thereby improving the public health of the populace.

## Introduction

Noise pollution is one of several environmental pollutions across the world. It can be described as the propagation of noise with a harmful impact on the physiological and psychological lives of humans or animals [1]. Noise or sound pollution is usually not studied compared with other forms of pollution such as air [2], [3], [4], water [5], soil [6], light and radioactive. The reason is that the adverse effects of other forms of pollution on humans are more pronounced. Notwithstanding, noise pollution remains a serious health concern in the study area (Ota, Nigeria) in particular and the entire planet [7], [8]. Some of the identified sources of noise pollution are loud music from concerts, religious buildings like churches and mosques, noise emitting generators [9], political rallies, road advertisement, traffic [10] and air transportation [11], sporting events, construction and industrial activities. In all the mentioned sources,

areas that have high risk of noise pollution are residential places near to major roads [12] and airports and manufacturing industries [13]; for example, small scale industries [14], [15], steel rolling industries [16], oil and gas industry [17], [18] and so on.

The health effects of noise pollution cannot be over-emphasised. This has prompted the World Health Organization (WHO) and the Federal Environment Protection Agency (FEPA) (Nigeria) to set standards and limits of allowable noise levels. Noise pollution occurs when it is observed that those standards are exceeded as seen in [19], [20].

The most common manifestation of noise pollution is hearing loss or impairment [21]. Hearing impairment is mostly classified as occupational hazards especially when the individual is affiliated with industry that propagates loud sound or noise. Moreover, several physiological and psychological effects of noise pollution exist. The combination of

noise and air pollution is associated with respiratory ailments, dizziness and tiredness in school children [22], [23]. In adults, noise pollution has been found to be associated with high blood pressure [24] and cognitive difficulties [25].

A look at the literature showed the abundance of evidence of the adverse effects of noise pollution on the general public health. The worsening situation of noise pollution is that it has not been upgraded to the level of the other forms of pollution. Also, recommendations suggested by several authors on the different strategies on tackling noise pollution has not been considered and implemented. However, noise pollution continues to impact negatively on fetal development [26], annoyance and anxiety [27], mental health crisis [28], sleep disturbance and insomnia [29], [30], cardiovascular disorders in pregnant women [31], cardiocerebrovascular diseases [32], type 2 diabetes incidence [33] and medically unexplained physical symptoms [34]. Other auditory and non-auditory effects of noise on health are myocardial infarction incidence [35], peptic ulcers [36] and disruption of communication and retentive capabilities in children [37].

This paper aims to study and analyse the noise pollution levels in major areas in Ota metropolis. A probability model which is capable of predicting the noise pollution level is also determined.

## Material and Methods

The dataset used in this research was gotten from [38]. It represents the noise level in 41 major locations in Ota metropolis, Nigeria. These major areas include industrial areas, commercial areas, passenger loading parks, busy roads and junctions. The readings were taken using the SLM (Sound Level Meter). Measurements were taken three different times of the day; morning (7 am to 9 am), afternoon (1 pm to 3 pm) and evening (6 pm to 8 pm). Particularly, the noise pollution level (NLP) was considered and analysed in this present research.




Figure 1: Summary report for morning measurements on LNP

## Analysis of Variance

Analysis of variance is conducted in this research to know if there is a significant difference between the effect of noise pollution level in the morning, afternoon and evening in Ota metropolis. The hypothesis tested is:

$H_0$ : The effects of the noise pollution level are the same for morning, afternoon and evening

Versus

$H_1$ : The effects of the noise pollution level are not the same for at least one of either morning, afternoon or evening.

The level of significance used is 0.05, and the null hypothesis is considered rejected if the p-value is less or equal to the level of significance. The structure of the ANOVA table is such as presented in Table 1.

Table 1: A typical example of a one-way ANOVA Table

Source of Variation	Degree of Freedom	Sum of Square	Mean Square	F-value
Factor	f-1	SSF	MSF = SSF/f-1	MSF/MSE
Error	n-f	SSE	MSE = SSE/n-f	
Total	n-1	SST		

where, 'f' is the number of factors which is 3 according to this research; morning, afternoon and evening. 'n' is the overall sample size.

## The goodness of Fit Test

The goodness of fit test is performed in this research to select the probability model that best fits the dataset. The Kolmogorov Smirnov (KS) test, the Anderson Darling (AD) test and Chi-square test are examples of the goodness of fit tests.




Figure 2: Summary report for afternoon measurements on LNP

The KS test was adopted in this research because it is the most popular and others might give similar results. The null hypothesis tests whether the data follow a specified distribution. If  $X_1, X_2, \dots, X_n$  represent ordered data points, the KS statistic is:

$$D \equiv \max_{1 \leq i \leq n} \left[ F(X_i) - \frac{i-1}{N}, \frac{i}{N} - F(X_i) \right]$$

where  $X_i$  are the ordered data and  $F(\cdot)$  is

the cumulative distribution function (cdf) of the continuous distribution tested.




Figure 3: Summary report for evening measurements on LNP




Figure 5: The 95% confidence interval (C.I) plot for the means

The result of the analysis of variance is presented in Table 3.

Table 3: Analysis of Variance (ANOVA) Table

Source	Degree of Freedom	Sum of Square	Mean Square	F-value	p-value
Factor	2	0.36	0.1805	0.00	0.997
Error	120	8585.85	71.5487		
Total	122	8586.21			

## Results

### Descriptive Analysis of the Dataset

The summary for the LNP measurements is provided in Figures 1 to 3 while the summary for the mean measurement across the 41 locations is provided in Figure 4.




Figure 4: Summary report for the mean measurements of LNP across all locations in Ota

The result in Table 3 shows that the generated p-value is 0.997 which is far greater than the level of significance (0.05). Hence, there is no enough evidence to reject the null hypothesis, and it can, therefore, be concluded that there is no significant difference in the means of the noise level measurements taken in the morning, afternoon and evening. This result is further confirmed by Turkey's post-hoc test which is summarized in Figure 6.




Figure 6: Summary of Turkey's post-hoc analysis

### Result for the Analysis of Variance

The analyses of the means of the various measurements are presented in Table 2.

Table 2: Analysis of the Means

Factor	N	Mean	Standard Deviation	95% Confidence Interval
LNP_Morning	41	90.78	7.89	(88.16, 93.39)
LNP_Afternoon	41	90.64	9.31	(88.03, 93.26)
LNP_Evening	41	90.72	8.11	(88.10, 93.34)

The 95% confidence interval (CI) plot for the means is displayed in Figure 5.

It can be observed in Figure 6 that all the intervals contained zero; this is an indication that there is no significant difference in the pair of each of the measurements considered.

### Fitting of Probability Models


To determine the appropriate probability model that describes the mean noise pollution level in Ota metropolis, Easyfit (trial version) software was used to select distribution with the best fit. The Kolmogorov-Smirnov (KS) test of goodness of fit was used to select the best model. The software fitted sixty distributions to the dataset, but the best five was

reported in this research. The result is presented in Table 4.

**Table 4: Fitted Distributions**

Distributions	KS Statistic	Rank
Log-Logistic (3P)	0.06236	1
Burr	0.06846	2
Hypersecant	0.07131	3
Logistic	0.08415	4
Johnson SU	0.08629	5

From Table 4, the best-fitted model is the three-parameter Log-logistic distribution; this selection/decision is based on the Kolmogorov Smirnov statistic. A graph showing the best distribution fitted to the dataset on mean noise pollution level is presented in Figure 7.



*Figure 7: Graph of log-logistic distribution on the histogram of the dataset*

In conclusion, further analyses of the noise pollution level in Ota metropolis has been provided in this research. The mean noise level in the morning was 90.78 which is higher than (though very close to) that of afternoon and evening with means 90.6 and 90.72 respectively. This is reasonable as more activities are expected during this time; pupils are going to school, workers going to the office, traffic at some junction and major bus stops. However, the analysis of variance result indicated that the time of the day (morning, afternoon and evening) have the same effect on the environment and populace. Also, the noise pollution level in Ota metropolis can be modelled using the log-logistic distribution as evident from the goodness of fit test. The model can now be used in predicting and managing noise pollution in that area. Furthermore, the model can be used in different geographical settings where noise pollution poses a perceived threat to the public health of the populace.

## References

- Oloruntoba EO, Ademola RA, Sridhar MKC, Agbola SA, Omokhodion FO, Ana GREE, Alabi RT. Urban environmental noise pollution and perceived health effects in Ibadan, Nigeria. *Afr J Biomed Res.* 2012; 15(2):77-84.
- Oguntunde PE, Odetunmbi OA, Adejumo AO. A study of probability models in monitoring environmental pollution in Nigeria. *J Prob Stat.* 2014; 2014: Article ID 864965. <https://doi.org/10.1155/2014/864965>
- Anake WU, Bayode FO, Omonhinmin CA, Williams AB. Ambient air pollution control using air pollution tolerance index and anticipated performance index of trees. *Int J Civil Eng Technol.* 2018; 9:417-425.
- Okokpuije K, Noma-Osaghae E, Modupe O, John S, Oluwatosin O. A smart air pollution monitoring system. *Int J Civil Engine Technol.* 2018; 9(9):799-809.
- Omole DO, Ogbiye AS, Longe EO, Adewumi IK, Elemile OO, Tenebe TI. Water quality checks on river atuwara, south-west Nigeria. *WIT Trans Ecol Environ.* 2018; 228:165-173. <https://doi.org/10.2495/WP180171>
- Oyeyemi KD, Aizebeokhai AP, Okagbue HI. Geostatistical exploration of dataset assessing the heavy metal contamination in Ewekoro limestone, Southwestern Nigeria. *Data in Brief.* 2017; 14:110-117. <https://doi.org/10.1016/j.dib.2017.07.041> PMID:28795088 PMCID:PMC5537382
- Egunjobi L. Urban environmental noise pollution in Nigeria. *Habitat Int.* 1986; 10(3):235-244. [https://doi.org/10.1016/0197-3975\(86\)90053-6](https://doi.org/10.1016/0197-3975(86)90053-6)
- Olokesusi F. An assessment of hotels in Abeokuta, Nigeria and its implications for tourists. *Int J Hospitality Magt.* 1990; 9(2):125-134. [https://doi.org/10.1016/0278-4319\(90\)90007-K](https://doi.org/10.1016/0278-4319(90)90007-K)
- Ibhadode O, Tenebe IT, Emenike PC, Adesina OS, Okougha AF, Aitanke FO. Assessment of noise-levels of generator-sets in seven cities of South-Southern Nigeria. *Afr J Sci Technol Innovat Develop.* 2018; 10(2):125-135. <https://doi.org/10.1080/20421338.2017.1400711>
- Oyedepo OJ, Ekom RI, Ajala KA. Analysis of traffic noise along oyemekun - oba-adesida road akure Ondo state Nigeria. *J Engine Sci Tech Rev.* 2013; 6(1):72-77. <https://doi.org/10.25103/jestr.061.14>
- Ibhadode O, Oyedepo OS, Ogunro AS, Azeta J, Solomon BO, Umanah II, Apeh ES, Ayoola AR. An Experimental-assessment of Human Exposure-levels to Aircraft Noise-hazards in the Neighbouring-environments of four Nigerian Airports, IOP Conf. Series: Mat Sci Engine. 2018; 413(1): Article number 012080. <https://doi.org/10.1088/1757-899X/413/1/012080>
- Asuquo U, Onuu M, Asuquo A. Effects of exposure to loud noise on the hearing of the residents of Calabar, Nigeria. *Canadian Acoustics.* 2012; 40(3):50-51.
- Bolaji BO, Olanipekun MU, Adekunle AA, Adeleke AE. An analysis of noise and its environmental burden on the example of Nigerian manufacturing companies. *J Cleaner Product.* 2018; 172:1800-1806. <https://doi.org/10.1016/j.jclepro.2017.12.007>
- Onuu MU, Akpan AO. Industrial noise in Nigeria: Measurements, analysis, dose and effects. *Building Acoustics.* 2006; 13(1):69-80. <https://doi.org/10.1260/135101006776324879>
- Oguntoke O, Odeshi, TA, Annegarn HJ. Assessment of noise emitted by vibrator-block factories and the impact on human health and urban environment in Nigeria. *Int J Appl Environ Sci.* 2012; 7(1):57-58.
- Ologe FE, Akande TM, Olajide TG. Occupational noise exposure and sensorineural hearing loss among workers of a steel rolling mill. *Euro Arch Oto-Rhino-Laryngol.* 2006; 263(7):618-621. <https://doi.org/10.1007/s00405-006-0043-9> PMID:16680467
- Abdulkareem AS, Odigure JO. Deterministic model for noise dispersion from gas flaring: A case study of Niger - Delta Area of Nigeria. *Chem Biochem Engine Quart.* 2006; 20(2):157-164.
- Aduloju AA, Okwechime I. Oil and human security challenges in the Nigeria's Niger delta. *Critique.* 2016; 44(4):505-525. <https://doi.org/10.1080/03017605.2016.1236495>
- Usikalu MR, Kolawole O. Assessment of noise pollution in selected locations in Ota, Nigeria. *Int J Mech Engine Technol.* 2018; 9(9):1212-1218.

20. Ogunsola OJ, Oluwole AF, Asubiojo OI, Durosinmi MA, Fatusi AO, Ruck W. Environmental impact of vehicular traffic in Nigeria: health aspects. *Sci Total Environ.* 1994; 146-147(C):111-116. [https://doi.org/10.1016/0048-9697\(94\)90226-7](https://doi.org/10.1016/0048-9697(94)90226-7)
21. Hinchcliffe R. Review: Global perspective of noise-induced hearing loss as exemplified by Nigeria. *J Audiolog Med.* 2002; 11(1):1-24.
22. Adetoun MB, Blangiardo M, Briggs DJ, Hansell AL. Traffic air pollution and other risk factors for respiratory illness in schoolchildren in the Niger-delta region of Nigeria. *Environ. Health Perspect.* 2011; 119(10):1478-1482. <https://doi.org/10.1289/ehp.1003099>
23. Shendell DG, Ana GREE, Brown GE, Sridhar MKC. Assessment of noise and associated health impacts at selected secondary schools in Ibadan, Nigeria. *J Environ Public Health Open.* 2009; 2009: Article number 739502. <https://doi.org/10.1155/2009/739502>
24. Ebare MN, Omuemu VO, Isah EC. Assessment of noise levels generated by music shops in an urban city in Nigeria. *Public Health.* 2011; 125(9):660-664. <https://doi.org/10.1016/j.puhe.2011.06.009> PMID:21875726
25. Ntui AI. Noise sources and levels at the University of Calabar Library, Calabar, Nigeria. *Afr J Libr Arch Info Sci.* 2009; 19(1):53-63.
26. Selander J, Rylander L, Albin M, Rosenhall U, Lewné M, Gustavsson P. Full-time exposure to occupational noise during pregnancy was associated with reduced birth weight in a nationwide cohort study of Swedish women. *Sci Total Environ.* 2019; 651:1137-1143. <https://doi.org/10.1016/j.scitotenv.2018.09.212> PMID:30360245
27. Paiva KM, Cardoso MR, Zannin PHT. Exposure to road traffic noise: Annoyance, perception and associated factors among Brazil's adult population. *Sci Total Environ.* 2019; 650:978-986. <https://doi.org/10.1016/j.scitotenv.2018.09.041> PMID:30308872
28. Freiberg A, Scheffer C, Girbig M, Murta VC, Seidler A. Health effects of wind turbines on humans in residential settings: Results of a scoping review. *Environ Research.* 2019; 169:446-463. <https://doi.org/10.1016/j.envres.2018.11.032> PMID:30530085
29. Eze IC, Foraster M, Schaffner E, Vienneau D, Héritier H, Pieren R, Thiesse L, Rudzik F, Rothe T, Pons M, Bettschart R, Schindler C, Cajochen C, Wunderli JM, Brink M, Röösli M, Probst-Hensch N. Transportation noise exposure, noise annoyance and respiratory health in adults: A repeated-measures study. *Environ Int.* 2018; 121:741-750. <https://doi.org/10.1016/j.envint.2018.10.006> PMID:30321849
30. Radun J, Hongisto V, Suokas M. Variables associated with wind turbine noise annoyance and sleep disturbance. *Build Environ.* 2019; 150:339-348. <https://doi.org/10.1016/j.buildenv.2018.12.039>
31. Sears CG, Braun JM, Ryan PH, Xu Y, Werner EF, Lanphear BP, Wellenius GA. The association of traffic-related air and noise pollution with maternal blood pressure and hypertensive disorders of pregnancy in the HOME study cohort. *Environ Int.* 2018; 121:574-581. <https://doi.org/10.1016/j.envint.2018.09.049> PMID:30300815
32. Oh M, Shin K, Kim K, Shin J. Influence of noise exposure on cardiocerebrovascular disease in Korea. *Sci Total Environ.* 2019; 651:1867-1876. <https://doi.org/10.1016/j.scitotenv.2018.10.081> PMID:30317174
33. Thiesse L, Rudzik F, Spiegel K, Leproult R, Pieren R, Wunderli JM, Foraster M, Héritier H, Eze IC, Meyer M, Vienneau D, Brink M, Probst-Hensch N, Röösli M, Cajochen C. Adverse impact of nocturnal transportation noise on glucose regulation in healthy young adults: Effect of different noise scenarios. *Environ Int.* 2018; 121:1011-1023. <https://doi.org/10.1016/j.envint.2018.05.036> PMID:30408889
34. Zock JP, Verheij R, Helbich M, Volker B, Spreeuwenberg P, Strak M, Janssen NAH, Dijst M, Groenewegen P. The impact of social capital, land use, air pollution and noise on individual morbidity in Dutch neighbourhoods. *Environ Int.* 2018; 121:453-460. <https://doi.org/10.1016/j.envint.2018.09.008> PMID:30273868
35. Bräuner EV, Jørgensen JT, Duun-Henriksen AK, Backalarz C, Laursen JE, Pedersen TH, Simonsen MK, Andersen Z. Long-term wind turbine noise exposure and incidence of myocardial infarction in the Danish nurse cohort. *Environ Int.* 2018; 121:794-802. <https://doi.org/10.1016/j.envint.2018.10.011> PMID:30336413
36. Min JY, Min KB. Cumulative exposure to nighttime environmental noise and the incidence of peptic ulcer. *Environ Int.* 2018; 121(Pt 2):1172-1178. <https://doi.org/10.1016/j.envint.2018.10.035>
37. Tesoriere G, Campisi T, Canale A, Severino A. The effects of urban traffic noise on children at kindergarten and primary school: A case study in Enna. *AIP Conf Proc.* 2018; 2040: Article number 140005. <https://doi.org/10.1063/1.5079194>
38. Oyedepo SO, Adeyemi GA, Fayomi OSI, Fagbemi OK, Solomon R, Adekeye T, Babalola OP, Akinyemi ML, Olawole OC, Joel ES, Nwanya SC. Dataset on noise level measurement in Ota metropolis, Nigeria. *Data in Brief.* 2019; 22:762-770. <https://doi.org/10.1016/j.dib.2018.12.049> PMID:30671520  
PMCID:PMC6330360