



Coupled transform method for time-space fractional Black-Scholes option pricing model



S.O. Edeki ^{a,*}, R.M. Jena ^b, S. Chakraverty ^b, D. Baleanu ^{c,d}

^a Department of Mathematics, Covenant University, Ota, Nigeria

^b Department of Mathematics, National Institute of Technology, Rourkela, India

^c Department of Mathematics, Faculty of Art and Sciences, Cankaya University, Balgat, 06530 Ankara, Turkey

^d Institute of Space Sciences, 077125 Magurele-Bucharest, Romania

Received 16 September 2019; revised 12 March 2020; accepted 21 August 2020

Available online 11 September 2020

KEYWORDS

Option pricing;
 Black Scholes model;
 RDTM;
 Fractional derivative;
 Analytical solutions

Abstract This paper presents analytical solutions of a time-space-fractional Black-Scholes model (TSFBSM) using a coupled technique referred to as Fractional Complex Transform (FCT) with the aid of a modified differential transform method. The nature of the derivatives is in the sense of Jumarie. The considered cases and applications show more consistency of the TSFBSM with an actual integer and fractional data when compared with the classical Black-Scholes model. The method is noted to be very effective, even with little knowledge of fractional calculus. Extension of this to multi-factor models formulated in terms of stochastic dynamics is highly recommended.

© 2020 The Authors. Published by Elsevier B.V. on behalf of Faculty of Engineering, Alexandria University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

From an industrial and academic point of view, the classical option pricing model referred to the Black-Scholes Model (BSM) is a partial differential equation (PDE) for the calculation of theoretical values of financial options based on some specific parameters such as current stock prices, expected volatility, time to expiration, strike prices, and expected dividends. The BSM remains a crucial financial tool for option valuation and pricing [1-5]. The relaxation of assumptions

governing the BSM most times leads to a non-linear version of the BSM or jump-diffusion models involving spikes [6-8]. In this work, the generalized form of the classical BSM for option pricing will be considered regarding time-space fractional order. The BSM for option pricing is of the form:

$$\frac{\partial \Lambda}{\partial t} + \frac{1}{2} S^2 \sigma^2 \frac{\partial^2 \Lambda}{\partial S^2} + rS \frac{\partial \Lambda}{\partial S} - r\Lambda = 0 \quad (1)$$

where $\Lambda = \Lambda(S, t)$ represents the option value, σ the volatility parameter, S , the price of underlying, r the interest rate (risk-free), and t as time. The associated payoff function, $p_{\Lambda}(S, t)$, and the price of expiration, E is denoted as:

$$p_{\Lambda}(S, t) = \pm c(S - E)^+ \quad (2)$$

for $c = \pm 1$ indicating European call and put option respectively, where $(S_*, 0)^+$ signifies the maximum between S_* and 0, σ as a parameter stands for the underlying asset ($S = S(t)$)

* Corresponding author.

E-mail addresses: soedeki@yahoo.com (S.O. Edeki), rajarama1994@gmail.com (R.M. Jena), sne_chak@yahoo.com (S. Chakraverty), dumitru@cankaya.edu.tr (D. Baleanu).

Peer review under responsibility of Faculty of Engineering, Alexandria University.

<https://doi.org/10.1016/j.aej.2020.08.031>

1110-0168 © 2020 The Authors. Published by Elsevier B.V. on behalf of Faculty of Engineering, Alexandria University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

volatility, r is the interest rate (in risk-free term), and T is the time at maturity. Suppose σ , the volatility parameter, and r the interest rate (risk-free) are constants, such that:

$$\left. \begin{aligned} S &= \zeta e^x, \\ \Lambda &= \Lambda(S, t) = \zeta \omega(x, \tau), \\ 2\tau &= (T - t)\sigma^2. \end{aligned} \right\} \tag{3}$$

Then, little algebra gives:

$$\left. \begin{aligned} \frac{\partial \Lambda}{\partial t} &= -\zeta \frac{\sigma^2}{2} \frac{\partial \omega}{\partial \tau}, \\ \frac{\partial \Lambda}{\partial S} &= \frac{\zeta}{S} \frac{\partial \omega}{\partial x}, \\ \frac{\partial^2 \Lambda}{\partial S^2} &= -\frac{\zeta}{S^2} \frac{\partial \omega}{\partial x} + \frac{\zeta}{S^2} \frac{\partial^2 \omega}{\partial x^2}. \end{aligned} \right\} \tag{4}$$

Thus, (4) in (1) gives:

$$\begin{aligned} -\zeta \frac{\sigma^2}{2} \frac{\partial \omega}{\partial \tau} + \frac{1}{2} S^2 \sigma^2 \left(-\frac{\zeta}{S^2} \frac{\partial \omega}{\partial x} + \frac{\zeta}{S^2} \frac{\partial^2 \omega}{\partial x^2} \right) + r S \left(\frac{\zeta}{S} \frac{\partial \omega}{\partial x} \right) - r(\zeta \omega(x, \tau)) &= 0. \\ \Rightarrow \left\{ \begin{aligned} \frac{\partial \omega}{\partial \tau} &= \frac{\partial^2 \omega}{\partial x^2} + (\kappa - 1) \frac{\partial \omega}{\partial x} - \kappa \omega, \quad \kappa = \frac{2r}{\sigma^2}, \\ \omega(x, 0) &= h(x). \end{aligned} \right. \end{aligned} \tag{5}$$

Equation (5) has received the attention of so many researchers in terms of effective semi-analytical numerical and analytical solution methods, which are of vital aid when handling differential models. Recently, in [9] the bond price model was extended to time-fractional order using conformable decomposition for analytical solutions. Edeki and Adinya [10] applied Fractional Complex Transform to a time-fractional one-factor Markovian model for bond pricing. Several methods of solution like variable separable method, direct integration, Laplace, Picard iteration method, differential transform method (DTM), Adomian decomposition method (ADM), Elzaki transform method (ETM), Sumudu decomposition method are applicable to PDEs [11-16].

Fractional calculus (FC) is used in different aspects of mathematics and applied sciences such as finance, biomedical, physics, and so on. FC helps in capturing the memory effect on the system.

With regard to FC and its applications, Morales-Delgado *et. al.*, [17] studied a non-integer order dynamics in relation to oxygen diffusion via the application of Laplace homotopy method. Zhang *et al.*, in [18] applied Laplace fractional homotopy perturbation method for the solution of heat conduction equations of non-homogeneous type involving fractional domains both in time- and space- derivatives. In [19], pathway fractional integer operator was considered with the aid of Mittag-Leffler functions. The authors in [20] presented some composition formulae for Marichev-Saigo-Maeda fractional operator, and obtained Laplace transforms of the corresponding composition formulae. In [21], some useful generalizations of fractional integro-differential operators on the basis of omega-parameter were introduced. Fractional kinetic equations based on hypergeometric functions were obtained in [22]. Damping characteristics of a fractional oscillator were considered in [23], while in [24], fractional oscillator model was used to describe damped vibrations. In [25], generalized fractional integral was utilized for obtaining Hermite-Hadamard-type inequalities for coordinated convex function. Several new properties of Mittag-Leffler function were recently obtained based on Konhauser and Laguerre polynomials [26], and [27], respectively. Jain *et. al.*, [28], considered certain

Hermite-Hadamard inequalities on the basis of convex logarithmic functions and derived new inequalities in terms of poly gamma functions.

For related researches on fractional models and solution methods with regard to detail and recent views, we make reference to [29-43] and the references therein.

The generalized version of (5) to time-space fractional-order to be considered takes the following form:

$$\left\{ \begin{aligned} \frac{\partial^\alpha \omega}{\partial \tau^\alpha} &= \frac{\partial^{2\gamma} \omega}{\partial x^{2\gamma}} + (\kappa - 1) \frac{\partial \omega}{\partial x} - \kappa \omega, \quad \alpha \in (0, 1], \quad 2\gamma \in (0, 2], \\ \omega(x, 0) &= h(x). \end{aligned} \right. \tag{6}$$

This paper, therefore, aims at providing approximate-analytical-solutions to the TSFBSM as contained in (6) for option pricing by applying the proposed semi-approximate technique known as Coupled Method. Similar PDEs can be considered by semi-analytical, numerical, and approximate methods in terms of solutions [44-49]. The fractional derivative in (6) is defined in the sense of Jumarie.

In financial mathematics, the fractional Black-Scholes model (FBSM) serves as a unique tool with respect to the modern theory of option pricing. The FBSM is a generalization of the classical BSM due to the modification(s) of the initial constancy assumptions. The exact/analytical solutions of which are difficult to obtain.

A lot of semi-analytical methods have been developed by numerical analysts for integer related mathematical and physical models. Meanwhile, modifications of some of these methods solve time-fractional models. It is, therefore, imperative to consider effective and reliable methods for space-fractional and time-space fractional models. Hence, the motivation for this research. The main feature of the proposed method of solution is hinged on the notion that it transforms the fractional model to its integer counterparts for ease of approach. This helps in employing some known classical semi-analytical methods such as the ADM, Differential Transform Method (DTM), Natural Decomposition Method (NDM), and so on.

2. Remarks on Jumarie’s fractional derivative

Reference is made to Fractional Derivative (JFD) in terms of Jumarie as a modified version Riemann-Liouville fractional derivatives [50,51]. The following is considered:

Consider $\Psi(z)$ as a real-valued function of z (continuous), and $D_z^\alpha \Psi = \frac{\partial^\alpha \Psi}{\partial z^\alpha}$ as JFD of $\Psi(z)$, whose order is α with respect to z . So,

$$D_z^\alpha \Psi = \begin{cases} \frac{1}{\Gamma(-\alpha)} \frac{d}{dz} \int_0^z (z - \zeta)^{-\alpha-1} (\Psi(\zeta) - \Psi(0)) d\zeta, & \alpha \in (-\infty, 0) \\ \frac{1}{\Gamma(1-\alpha)} \frac{d}{dz} \int_0^z (z - \zeta)^{-\alpha} (\Psi(\zeta) - \Psi(0)) d\zeta, & \alpha \in (0, 1) \\ (\Psi^{(\alpha-\eta)}(z))^{(\eta)}, & \alpha \in [\eta, \eta + 1), \quad \eta \geq 1 \end{cases} \tag{7}$$

where $\Gamma(\cdot)$ represents a gamma function. The main features of JFD [51] are as follows:

- (i). $D_z^\alpha c = 0, \alpha > 0$, for a constant c
- (ii). $D_z^\alpha (c\Psi(z)) = cD_z^\alpha \Psi(z), \alpha > 0$,
- (iii). $D_z^\alpha z^\beta = \begin{cases} \frac{\Gamma(\beta+1)}{\Gamma(\beta+1-\alpha)} z^{\beta-\alpha}, & \text{for } \beta \neq 1 - \alpha. \end{cases}$
- (iv). $D_z^\alpha (\Psi_1(z)\Psi_2(z)) = D_z^\alpha \Psi_1(z)(\Psi_2(z)) + \Psi_1(z)D_z^\alpha \Psi_2(z),$
- (v). $D_z^\alpha (\Psi(z)g(z)) = D_z^\alpha \Psi \cdot D_z^\alpha g(z).$

The properties in (i) through (v) are basic in applications.

3. The Reduced differential Transform [30]

Let $\omega(x, t)$ be a continuously differentiable and analytic function on a given domain, D , so the differential transform of $\omega(x, t)$ is expressed as:

$$\Omega_k(x) = \frac{1}{k!} \left[\frac{\partial^k \omega(x, t)}{\partial t^k} \right]_{t=0} \tag{8}$$

where $\omega(x, t)$ and $\Omega_k(x)$ are referred to as the original and the transformed functions in that order. Hence, the differential inverse transform (DIT) of $\Omega_k(x)$ is given as:

$$\omega(x, t) = \sum_{k=0}^{\infty} \Omega_k(x) t^k \tag{9}$$

3.1. DTM and its basic properties

D1: If $\omega(x, t) = \alpha q(x, t) \pm \beta p(x, t)$, then $\Omega_k(x) = \alpha Q_k(x) \pm \beta P_k(x)$.

D2: If $\omega(x, t) = \frac{c \partial^m q(x, t)}{\partial t^m}$, $m \in N$, then $\Omega_k(x) = \frac{c(k+m)!}{k!} Q_{k+m}(x)$.

D3: If $\omega(x_*, t) = \frac{m(x_*) \partial^{\eta} q(x_*, t)}{\partial x_*^{\eta}}$, $\eta \in N$, then $\Omega_k(x_*) = \frac{M(x_*) \partial^{\eta} Q_k(x_*)}{\partial x_*^{\eta}}$, $\eta \in N$.

D4: If $\omega(x, t) = p(x, t) q(x, t)$, then $\Omega_k(x) = \sum_{\eta=0}^k P_{\eta}(x) Q_{k-\eta}(x)$.

D5: If $\omega(x, t) = x^{n_1} t^{n_2}$, then $\Omega_c = x^{n_1} \delta(c - n_2)$, $\delta(c) = \begin{cases} 0, & c \neq 0, \\ 1, & c = 0. \end{cases}$

3.2. The fractional Complex Transform

The general fractional differential equation of the following form is considered:

$$h(v, D_t^{\alpha} v, D_x^{\beta} v, D_y^{\gamma} v, D_z^{\delta} v) = 0, \quad v = v(t, x, y, z) \tag{10}$$

while the FCT as considered in [30,31], and the references therein is defined as follows:

$$T = \frac{at^{\alpha}}{\Gamma(1+\alpha)}, \quad \alpha \in (0, 1]. \tag{11}$$

Thus, using (iii) in section 2, we have:

$$D_z^{\delta} z^{\beta} = \frac{\Gamma(\beta+1)}{\Gamma(\beta+1-\alpha)} z^{\beta-\alpha}, \quad 0 < \alpha \leq \beta,$$

$$\therefore D_t^{\alpha} T = \frac{a}{\Gamma(1+\alpha)} \left[\frac{\Gamma(\alpha+1)}{\Gamma(\alpha-\alpha+1)} \right] t^{\alpha-\alpha} = a. \tag{12}$$

Hence,

$$\begin{cases} D_t^{\alpha} \omega = D_t^{\alpha} \omega(T(\tau)) = D_T^1 \omega \cdot D_t^{\alpha} T = a \frac{\partial \omega}{\partial T}, \\ D_x^{\beta} \omega = D_x^{\beta} \omega(X(t)) = D_X^1 \omega \cdot D_X^{\beta} X = b \frac{\partial \omega}{\partial X}. \end{cases} \tag{13}$$

4. Applications

In this section, the proposed FCT is applied to time-space fractional Black-Scholes models (TSFBSMs) as follows:

Case I- Consider the following form of TSFBSM:

$$\begin{cases} \frac{\partial^2 \omega}{\partial \tau^2} = \frac{\partial^2 \omega}{\partial x^2} + (\kappa - 1) \frac{\partial \omega}{\partial x} - \kappa \omega, \quad \alpha \in (0, 1], \quad 2\gamma \in (0, 2], \\ \omega(x, 0) = (\exp(x) - 1)^+. \end{cases} \tag{14}$$

It is remarked from FCT, that $T = \frac{a\tau^{\alpha}}{\Gamma(1+\alpha)}$, $X = \frac{ax^{\gamma}}{\Gamma(1+\gamma)}$, which according to section 3 gives

$$\frac{\partial^{\alpha} \omega}{\partial \tau^{\alpha}} = \frac{\partial \omega}{\partial T}, \quad \frac{\partial^{\gamma} \omega}{\partial x^{\gamma}} = \frac{\partial \omega}{\partial X}, \quad a = b = 1.$$

Hence, (12) becomes:

$$\begin{cases} \frac{\partial \omega}{\partial T} = \frac{\partial^2 \omega}{\partial X^2} + (\kappa - 1) \frac{\partial \omega}{\partial X} - \kappa \omega, \\ \omega(X, 0) = \max(e^X - 1, 0). \end{cases} \tag{15}$$

Applying the RDTM to (15) gives:

$$\begin{cases} W_{k+1} = \frac{1}{(1+k)} \left(W''_{X,k} + (\kappa - 1) W'_{X,k} - \kappa W_k \right), \quad k \geq 0, \\ W(X, 0) = \max(e^X - 1, 0), \end{cases} \tag{16}$$

where the prime notations denote derivatives w.r.t. variable X . As a result, the recursive relation in (16) yields:

$$\begin{cases} W_0 = \max\{\exp(X) - 1, 0\}, \\ W_1 = \kappa \{-\max(\exp(X) - 1, 0) + \max(\exp(X), 0)\}, \\ W_2 = \frac{-\kappa^2}{2} \{-\max(\exp(X) - 1, 0) + \max(\exp(X), 0)\}, \\ W_3 = \frac{\kappa^3}{6} \{-\max(\exp(X) - 1, 0) + \max(\exp(X), 0)\}, \\ \vdots \\ W_p = \frac{(-1)^{p+1} (\kappa)^p}{p!} \left\{ \max_{-\max(0, -\exp(X))}^{(0, \exp(X))} \right\}, \quad p = 1, 2, \dots \end{cases} \tag{17}$$

Hence,

$$\begin{aligned} \omega(X, T) &= \sum_{m=0}^{\infty} W_m T^m = W_0 + \sum_{m=1}^{\infty} W_m T^m \\ &= \max(\exp(X) - 1, 0) + \sum_{m=1}^{\infty} (-1)^{m+1} \frac{(\kappa T)^m}{m!} \left\{ \begin{array}{l} \max(\exp(X), 0) \\ -\max(\exp(X) - 1, 0) \end{array} \right\} \end{aligned} \tag{18}$$

Therefore, the solution of (14) is as follows:

$$\omega(x, \tau) = \left\{ \begin{array}{l} \max\left(\exp\left(\frac{x^{\gamma}}{\Gamma(1+\gamma)}\right) - 1, 0\right) \\ + \sum_{m=1}^{\infty} (-1)^{m+1} \frac{1}{m!} \left\{ \frac{\kappa \tau^{\alpha}}{\Gamma(1+\alpha)} \right\}^m \left\{ \begin{array}{l} \max\left(\exp\left(\frac{x^{\gamma}}{\Gamma(1+\gamma)}\right), 0\right) \\ -\max\left(\exp\left(\frac{x^{\gamma}}{\Gamma(1+\gamma)}\right) - 1, 0\right) \end{array} \right\} \end{array} \right\} \tag{19}$$

In (19), the solution to the TSFBSM for option pricing (14) is provided. At $\alpha = \gamma = 1$, the solutions obtained in [52,53] serve as particular cases to this present work.

Case II- Consider the following TSFBSM of the form:

$$\begin{cases} \frac{\partial^2 \omega}{\partial \tau^2} = \omega - \left(\frac{x}{2} \frac{\partial \omega}{\partial x} + x^2 \frac{\partial^2 \omega}{\partial x^2} \right), \quad \alpha \in (0, 1], \quad 2\gamma \in (0, 2], \\ \omega(x, 0) = \max(x^3, 0) = \begin{cases} x^3, & x > 0 \\ 0, & x \leq 0. \end{cases} \end{cases} \tag{20}$$

This coincides with *problem 1* of [54] for $\gamma = 1$ and [55] for $\alpha = \gamma = 1$ as particular cases of this present work.




Fig. 1 Case I Solution plot for $\alpha = \gamma = 1$




Fig. 3 Case I Solution plot for $\alpha = 1, \gamma = 0.5$




Fig. 2 Case I Solution plot for $\alpha = 0.5, \gamma = 1$




Fig. 4 Case I Solution plot for $\alpha = 0.75, \gamma = 0.75$

Thus, following the FCT as in the previous case with:

$$\frac{\partial^x \omega}{\partial \tau^x} = \frac{\partial \omega}{\partial T}, \quad \frac{\partial^y \omega}{\partial x^y} = \frac{\partial \omega}{\partial X}, \quad a = b = 1,$$

reforms (20) to:

$$\begin{cases} \frac{\partial \omega}{\partial T} = \omega - \left(\frac{X}{2} \frac{\partial \omega}{\partial X} + X^2 \frac{\partial^2 \omega}{\partial X^2} \right), \\ \omega(X, 0) = \max(X^3, 0). \end{cases} \quad (21)$$

Applying the RDTM to (21) gives:

$$\begin{cases} W_{k+1} = \frac{1}{(1+k)} \left\{ W_k - \left(\frac{X}{2} W'_{X,k} + X^2 W''_{X,k} \right) \right\}, \quad k \geq 0, \\ W(X, 0) = \max(X^3, 0). \end{cases} \quad (22)$$

As a result, the following are yielded:

$$\begin{cases} W_0 = X^3, \quad W_1 = (-6.5)X^3, \\ W_2 = \frac{(-6.5)^2}{2!} X^3, \quad W_3 = \frac{(-6.5)^3}{3!} X^3, \\ W_4 = \frac{(-6.5)^4}{4!} X^3, \\ \vdots \\ W_p = \frac{(-6.5)^p}{p!} X^3, \quad p = 0, 1, 2, \dots \end{cases} \quad (23)$$




Fig. 5 Case I Solution plot for $\alpha = 0.95, \gamma = 0.25$




Fig. 7 Case II Solution plot for $\alpha = 0.5, \gamma = 1$




Fig. 6 Case II Solution plot for $\alpha = \gamma = 1$




Fig. 8 Case II Solution plot for $\alpha = 1, \gamma = 0.5$

Therefore,

$$\begin{aligned} \omega(X, T) &= \sum_{m=0}^{\infty} W_m T^m = W_0 + \sum_{m=1}^{\infty} W_m T^m \\ &= X^3 + \sum_{m=1}^{\infty} \left(\frac{(-6.5)^m}{m!} X^3 \right) T^m \\ &= X^3 \sum_{m=0}^{\infty} \left(\frac{(-6.5T)^m}{m!} \right) = X^3 \exp(-6.5T). \end{aligned} \tag{24}$$

Hence, the solution of (20) is given as:

$$\omega(x, \tau) = \left(\frac{x^\gamma}{\Gamma(1+\gamma)} \right)^3 \exp\left(\frac{-6.5\tau^\alpha}{\Gamma(1+\alpha)} \right). \tag{25}$$

Note: At $\alpha = \gamma = 1$, the exact solution [54] is:

$$\omega(x, \tau) = x^3 \exp(-6.5\tau). \tag{26}$$

In (26), the solution to the TSFBSM for option pricing (20) is provided. At $\alpha = \gamma = 1$, the solutions obtained in [32,54,56] serve as particular cases to this present work.

4.1. Notes on the numerical solution

It is noteworthy here, that Fig. 1 shows the solution graphic of the model in case I for $\alpha = \gamma = 1$, which implies the non-fractional (classical integer) case of the model. Fig. 2 and Fig. 3 denote plots for purely time and purely space fractional cases, respectively. Meanwhile, Fig. 4 and Fig. 5 represent the solution plots of the model for time-space fractional cases.




Fig. 9 Case II Solution plot for $\alpha = 0.75, \gamma = 0.75$




Fig. 10 Case II Solution plot for $\alpha = 0.95, \gamma = 0.25$

Similarly, we have in Figs. 6-10 for case II. Tables 1 and 2 show the comparison of present results with the exact solution at a particular value of $\gamma = 1$.

5. Concluding remarks

This paper presented a solution method referred to as Coupled Transform Method with combined features of Fractional Complex Transform and Reduced Differential Transform Method. This proposed method is applied for an approximate-analytical-solution of a time-space fractional Black-Scholes model (TSFBSM) for option pricing. It is worthy of remark that this proposed method does not necessarily require a complete knowledge of fractional calculus while com-

Table 1 Comparison of the present solution with the exact solution of Case I.

τ	x	$\alpha = 1.0, \kappa = 2, \text{ and } \gamma = 1$	
		Present solution	Exact solution
0.2	0.00	0.3296799	0.3296799
	0.25	0.6137053	0.6137053
	0.50	0.9784012	0.9784012
	0.75	1.4466799	1.4466799
	1.00	2.0479617	2.0479617
0.4	0.00	0.5506710	0.5506710
	0.25	0.8346964	0.8346964
	0.50	1.1993923	1.1993923
	0.75	1.6676710	1.6676710
	1.00	2.2689528	2.2689528
0.6	0.00	0.6988057	0.6988057
	0.25	0.9828312	0.9828312
	0.50	1.3475270	1.3475270
	0.75	1.8158058	1.8158058
	1.00	2.4170876	2.4170876

Table 2 Comparison of the present solution with the exact solution of Case II.

τ	x	$\alpha = 1.0 \text{ and } \gamma = 1$	
		Present solution	Exact solution
0.2	0.00	0.000000000000	0.000000000000
	0.25	0.4258309266E-2	0.4258309266E-2
	0.50	0.3406647413E-1	0.3406647413E-1
	0.75	0.114974350200	0.114974350200
	1.00	0.272531793000	0.272531793000
0.4	0.00	0.000000000000	0.000000000000
	0.25	0.1160524659E-2	0.1160524659E-2
	0.50	0.9284197273E-2	0.9284197273E-2
	0.75	0.3133416580E-1	0.3133416580E-1
	1.00	0.7427357819E-1	0.7427357819E-1
0.6	0.00	0.000000000000	0.000000000000
	0.25	0.3162798663E-3	0.3162798663E-3
	0.50	0.2530238930E-2	0.2530238930E-2
	0.75	0.8539556390E-2	0.8539556390E-2
	1.00	0.2024191144E-1	0.2024191144E-1

puting the solutions of fractional models, yet the level of accuracy is highly maintained.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The first author thanks Covenant University for the provision of a good working environment. Also, the second-named author thanks the Department of Science and Technology of the Government of India regarding the provision of INSPIRE Fellowship (IF170207) for this research.

References

- [1] J.C. Hull, *Options, Futures, and other derivatives*, Prentice-Hall, New Jersey, 1999.
- [2] F. Black, M. Scholes, The pricing of options and corporate liabilities, *Journal of Political Economy* 81 (1973) 637–654.
- [3] R.C. Merton, Theory of rational options pricing, *Bell J. Econ. Manag. Sci.* 4 (1) (1973) 141–183.
- [4] S.O. Edeki, E.A. Owoloko, O.O. Ugbebor, The Modified Black-Scholes Model via Constant Elasticity of Variance for Stock Options Valuation, *AIP Conf. Proc.* 1705 (2016), <https://doi.org/10.1063/1.4940289> 020041.
- [5] O. González-Gaxiola, J.R. de Chávez, J.A. Santiago, A Nonlinear Option Pricing Model Through the Adomian Decomposition Method, *Int. J. Appl. Comput. Math* (2015), <https://doi.org/10.1007/s40819-015-0070-6>.
- [6] S. Prabhakaran, Construction of Pde Black - Scholes with Jump-Diffusion Models, *International Research Journal of Finance and Economics* 168 (2018) 48–67.
- [7] A. Ciancio, Analysis of time series with wavelets, *Int. J. Wavelets Multiresolut. Inf. Process.* 5 (2) (2007) 241–256.
- [8] L. Boen, K. J., In 't Hout, Operator splitting schemes for American options under the two-asset Merton jump-diffusion model, *Applied Numerical Mathematics*, 153, (2020), 114-131.
- [9] S.O. Edeki, I. Adinya, G.O. Akinlabi, O.P. Ogundile, Conformable decomposition for analytical solutions of a time-fractional one-factor Markovian model for bond pricing, *Applied Mathematics & Information Sciences* 13 (4) (2019) 1–6.
- [10] S.O. Edeki, I. Adinya, Coupled transform for approximate-analytical solutions of a time-fractional one-factor markovian bond pricing model, *WSEAS Transactions on Systems and Control* 14 (2019) 202–208.
- [11] F. Mehrdoust, M. Mirzazadeh, On Analytical Solution of the Black-Scholes Equation by the First Integral Method, *U.P.B. Sci. Bull., Series A* 76 (4) (2014) 85–90.
- [12] V. Gülkaç, The homotopy perturbation method for the Black-Scholes equation, *J. Stat. Comput. Simul.* 80 (12) (2010) 1349–1354.
- [13] J. Esekun, S. Onyango, N.O. Ongati, Analytic solution of a nonlinear Black-Scholes partial differential equation, *International Journal of Pure and Applied Mathematics* 61 (2) (2010) 219–230.
- [14] J. Ankudinova, M. Ehrhardt, On the numerical solution of nonlinear Black-Scholes equations, *Comput. Math. Appl.* 56 (2008) 799–812.
- [15] S.O. Edeki, O.O. Ugbebor, and E.A. Owoloko, He's Polynomials for Analytical Solutions of the Black-Scholes Pricing Model for Stock Option Valuation, *Proceedings of the World Congress on Engineering*, London, UK, (2016).
- [16] J.H. He, Homotopy perturbation method: A new nonlinear analytical technique, *Appl. Math. Comput.* 135 (2003) 73–79.
- [17] V.F. Morales-Delgado, J.F. Gómez-Aguilar, Khaled M. Saad, Muhammad Altaf Khan, P. Agarwal, Analytic solution for oxygen diffusion from capillary to tissues involving external force effects: A fractional calculus approach, *Physica A* 523 (2019) 48–65.
- [18] Y. Zhang, C. Cattani, X.J. Yang, Local fractional homotopy perturbation method for solving non-homogeneous heat conduction equations in fractal domains, *Entropy* 17 (10) (2015) 6753–6764.
- [19] S. Jain, P. Agarwal, A. Kilicman, Pathway Fractional Integral Operator Associated with 3m-Parametric Mittag-Leffler Functions, *International Journal of Applied and Computational Mathematics* 4 (2018) 115.
- [20] S. Jain, P. Agarwal, I. O. Kıymaz and A. Cetinkaya, Some composition formulae for the MSM fractional integral operator with the multi-index Mittag-Leffler functions, *AIP Conference Proceedings* 1926 (1), 020020, (2018).
- [21] P. Agarwal, A.M. Jerbashian, J.E. Restrepo, Weighted Integral Inequalities in Terms of Omega-Fractional Integro-Differentiation, in: P. Agarwal, S. Dragomir, M. Jleli, B. Samet (Eds.), *Advances in Mathematical Inequalities and Applications*, Trends in Mathematics, Birkhäuser, Singapore, 2018.
- [22] K.B. Kachhia, P. Agarwal, J.C. Prajapati, Certain Image Formulae and Fractional Kinetic Equations Involving Extended Hypergeometric Functions, in: M. Ruzhansky, Y. Cho, P. Agarwal, I. Area (Eds.), *Advances in Real and Complex Analysis with Applications*, Trends in Mathematics, Birkhäuser, Singapore, 2017.
- [23] B.N. Narahari Achar, J.W. Hanneken, T. Clarke, Damping characteristics of a fractional oscillator, *Physica A* 339 (2004) 311–319, <https://doi.org/10.1016/j.physa.2004.03.030>.
- [24] S. Rekhviashvili, A. Pskhu, P. Agarwal, S. Jain, Application of the fractional oscillator model to describe damped vibrations, *Turkish Journal of Physics* 43 (3) (2019) 236–242.
- [25] H. Budak, P. Agarwal, On Hermite–Hadamard-Type Inequalities for Coordinated Convex Mappings Utilizing Generalized Fractional Integrals. In: Agarwal P., Baleanu D., Chen Y., Momani S., Machado J. (eds) *Fractional Calculus. ICFDA 2018*. Springer Proceedings in Mathematics & Statistics, 303 (2019), Springer, Singapore.
- [26] J.C. Prajapati., N.K. Ajudia , S. Jain, A. Goswami, P. Agarwal, Certain Properties of Konhauser Polynomial via Generalized Mittag-Leffler Function. In: Agarwal P., Baleanu D., Chen Y., Momani S., Machado J. (eds) *Fractional Calculus. ICFDA 2018*. Springer Proceedings in Mathematics & Statistics, 303, (2019), Springer, Singapore.
- [27] P. Agarwal, M. Chand, S. Rani, T.M. Rassias, Certain Feynman type integrals involving generalized k-Mittag-Leffler function and general class of polynomials, *Applicable Analysis and Discrete Mathematics* (2019) 31.
- [28] S. Jain, K. Mehrez, D. Baleanu, P. Agarwal, Certain Hermite Hadamard inequalities for logarithmically convex functions with applications, *Mathematics* 7 (2) (2019) 163.
- [29] S.O. Edeki, G.O. Akinlabi, Coupled method for solving time-fractional navier stokes equation, *International Journal of Circuits, Systems and Signal Processing* 12 (2017) 27–34.
- [30] S.O. Edeki, G.O. Akinlabi, S.A. Adeosun, Analytic and Numerical Solutions of Time-Fractional Linear Schrödinger Equation, *Comm Math Appl* 7 (1) (2016) 1–10.
- [31] S.O. Edeki, O.P. Ogundile, B. Osoba, G.A. Adeyemi, F.O. Egara, S.A. Ejoh, Coupled FCT-HP for Analytical Solutions of the Generalized Time-Fractional Newell-Whitehead-Segel Equation, *WSEAS Transactions on Systems and Control* 13 (2018) 266–274.
- [32] S.O. Edeki, O.O. Ugbebor, E.A. Owoloko, Analytical Solution of the Time-fractional Order Black-Scholes Model for Stock Option Valuation on No Dividend Yield Basis, *IAENG International Journal of Applied Mathematics* 47 (4) (2017) 407–416.
- [33] S.O. Edeki, G.O. Akinlabi, Zhou Method for the Solutions of System of Proportional Delay Differential Equations, *MATEC Web of Conferences*, 125 (2017), Article number 02001.
- [34] A.A Yildirim, Application of the Homotopy perturbation method for the Fokker-Planck equation, *International Journal for Numerical Methods in Biomedical Engineering* 26 (2010) 1144–1154.
- [35] Owolabi K., Atangana A., On the formulation of Adams-Bashforth scheme with Atangana-Baleanu-Caputo fractional derivative to model chaotic problems, *Chaos* 29(2) (2019), 023111. <https://doi.org/10.1063/1.5085490>.

- [36] E. Ozbilge, A. Demir, An inverse source problem in time-space fractional differential equations, *Applied Mathematics and Information Sciences* 12 (3) (2018) 587–591.
- [37] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier Science B.V, Amsterdam, 2006.
- [38] I. Podlubny, *Fractional Differential Equation*, Academic Press, San Diego, 1999.
- [39] D. Baleanu, J.A.T. Machado, A.C. Luo, *Fractional Dynamics and Control*, Springer, Berlin, Germany, 2012.
- [40] R.M. Jena, S. Chakraverty, Solving time-fractional Navier-Stokes equations using homotopy perturbation Elzaki transform, *SN Applied Sciences* 1 (1) (2019) 16.
- [41] R.M. Jena, S. Chakraverty, A new iterative method based solution for fractional Black-Scholes option pricing equations (BSOPE), *SN Applied Sciences* 1 (2019) 95.
- [42] R.M. Jena, S. Chakraverty, Residual Power Series Method for Solving Time-fractional Model of Vibration Equation of Large Membranes, *J. Appl. Comput. Mech.* 5 (4) (2019) 603–615.
- [43] R.M. Jena, S. Chakraverty, D. Baleanu, On New Solutions of Time-Fractional Wave Equations Arising in Shallow Water Wave Propagation, *Mathematics* 7 (2019) 722.
- [44] D. Baleanu, K. Diethelm, E. Scalas, J.J. Trujillo, *Fractional Calculus: Models and Numerical Methods*; World Scientific Publishing Company: Boston, MA, USA, 2012.
- [45] S. Javeed, D. Baleanu, A. Waheed, M.S. Khan, H. Affan, Analysis of Homotopy Perturbation Method for Solving Fractional Order Differential Equations, *Mathematics* 7 (2019) 40.
- [46] S.O. Edeki, T. Motsepa, C.M. Khalique, G.O. Akinlabi, The Greek parameters of a continuous arithmetic Asian option pricing model via Laplace Adomian decomposition method, *Open Physics* 16 (2018) 780–785.
- [47] Z. A. Elshegmani, R. R. Ahmad Analytical solution for an arithmetic Asian option using mellin transforms. *Int. Journal of Math. Analysis*, 5 (26), (2011), 1259-1265.
- [48] C. Cattani, T.A. Sulaiman, H.M. Baskonus, H. Bulut, On the soliton solutions to the Nizhnik-Novikov-Veselov and the Drinfeld-Sokolov systems, *Opt. Quant. Electron.* 50 (3) (2018) 138.
- [49] G.O. Akinlabi, S.O. Edeki, The Solution of Initial-value Wave-like Models via Perturbation Iteration Transform Method, *Lecture Notes in Engineering and Computer Science* 2228 (2017) 1015–1018.
- [50] T.M. Elzaki, The New Integral Transform “Elzaki Transform”, *Global Journal of pure and applied mathematics*, ISSN 0973–1768 6 (1) (2011) 57–64.
- [51] C. Cattani, Haar wavelet-based technique for sharp jumps classification, *Math. Comput. Modell.* 39 (2–3) (2004) 255–278.
- [52] G. Jumarie, Fractional partial differential equations and modified Riemann- Liouville derivative new methods for solution, *J. Appl. Math. Comput.* 24 (1–2) (2007) 31–48.
- [53] G. Jumarie, Modified Riemann-Liouville Derivative and Fractional Taylor series of Non-differentiable Functions Further Results, *Comput. Math. Appl.* 51 (9–10) (2006) 1367–1376.
- [54] S.O. Edeki, O.O. Ugbebor, E.A. Owoloko, Analytical Solutions of the Black-Scholes Pricing Model for European Option Valuation via a Projected Differential Transformation Method, *Entropy* 17 (11) (2015) 7510–7521.
- [55] S. Kumar, A. Yildirim, Y. Khan, H. Jafari, K. Sayevand, L. Wey, Analytical solution of fractional Black-Scholes, European option pricing equation by using Laplace transform, *J. of Fractional Calculus and Applications* 2 (2012) 1–9.
- [56] T. Allahviranloo, Sh.S. Behzadi, The use of iterative methods for solving Black-Scholes equation, *International Journal of Industrial Mathematics* 5 (1) (2013) 1–11.