

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/334466355>

Conformable Decomposition for Analytical Solutions of a Time-Fractional One-Factor Markovian Model for Bond Pricing

Article · January 2019

CITATIONS

0

READS

94

4 authors:



S.O. Edeki

Covenant University Ota Ogun State, Nigeria

127 PUBLICATIONS 734 CITATIONS

[SEE PROFILE](#)



Ini Adinya

University of Ibadan

12 PUBLICATIONS 32 CITATIONS

[SEE PROFILE](#)



G.O. Akinlabi

Covenant University Ota Ogun State, Nigeria

40 PUBLICATIONS 317 CITATIONS

[SEE PROFILE](#)



Opeyemi Paul Ogundile

Covenant University Ota Ogun State, Nigeria

36 PUBLICATIONS 102 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Karhunen-Loeve expansion of stochastic dynamics with applications to a certain class of financial model [View project](#)



Thesis [View project](#)

Conformable Decomposition for Analytical Solutions of a Time-Fractional One-Factor Markovian Model for Bond Pricing

S. O. Edeki^{1,*}, I. Adinya², G. O. Akinlabi¹ and O. P. Ogundile¹

¹Department of Mathematics, Covenant University, Ota, Nigeria

²Department of Mathematics, University of Ibadan, Nigeria

Received: 3 Oct. 2018, Revised: 16 Feb. 2019, Accepted: 23 May 2019

Published online: 1 Jul. 2019

Abstract: In financial and option pricing setting, one-factor model denotes the notion that there exists one Wiener process in the definition of the short-rate process indicating one source of randomness. In this paper, approximate-analytical solution of a time-fractional one-factor Markovian model for bond pricing is considered using the approach of conformable decomposition. The method is a modified version of Adomian decomposition coupled with fractional derivative defined in conformable sense. Illustrative examples are presented in order to clarify the effectiveness of the proposed solution method, and the solutions are presented graphically based on some financial parameters at different values of the time-fractional order. This approach can be extended to multi-factor models formulated in terms of stochastic dynamics.

Keywords: Option pricing, Black Scholes model, Adomian decomposition, conformable fractional derivative, analytical solutions

1 Introduction

In any financial setting, interest rate implies a certain rate (amount) that is charged upon the use of financial item, mainly money [1]. In financial market, interest rates are non-tradable assets but are derived from the prices of tradable assets such as bonds and swaps. An Interest Rate Model (IRM) describes the evolution of the interest rates in relation to their dependency at maturity known as Term Structure of Interest Rates (TSIR) [2, 3]. Meanwhile, bond prices are popularly modeled using short-rate models [1, 4]. It is basically assumed that the short rate, ξ is governed by the general stochastic differential equation (SDE):

$$d\xi = \mu(\xi, t)dt + \sigma(\xi, t)dW \quad (1)$$

where $\mu(\xi, t)$, $\sigma(\xi, t)$, and $W = W(t)$ are trend process, random fluctuation determinant function, and Wiener process respectively. Equation (1) yields Markovian models if $\mu(\xi, t) = \mu(\xi)$ and $\sigma(\xi, t) = \sigma(\xi)$; meaning that μ and σ are strictly functions of variable ξ (not depending on t at all). We speak of one-factor models or multi-factor models if $\xi = X$ in (1) is a scalar or a vector

respectively [5]. This work considers one-factor model (denoting the existence of one Wiener process-for one source of randomness).

The choice of the volatility function with respect to (w.r.t.) correctness has been a topic of concern. Chen, Karolyi, Longstaff, and Sanders proposed a general Short Rate Model (SRM) defined in terms of single stochastic dynamics of the form:

$$d\xi = (\beta_1 + \beta_2\xi)dt + \sigma\xi^\phi dW \quad (2)$$

known as CKLS model; where β_1 , β_2 , ϕ are constants, and $W = W(t)$ is a Wiener process [6]. In this regard, related references include those of [7-16]. By considering only corresponding Markov models, and taking $\tau = T - t$ (for convenient) to denote the remaining time to maturity, it is therefore claimed that the bond price, $P(\xi, \tau)$ solves the parabolic partial differential model evolving from (2) expressed in the form:

$$\begin{cases} \frac{\partial P}{\partial \tau} = \frac{1}{2}\sigma^2\xi^{2\phi}\frac{\partial^2 P}{\partial \xi^2} + (\beta_1 + \beta_2\xi)\frac{\partial P}{\partial \xi} - \xi P \\ P(\xi, 0) = h(\xi) \end{cases} \quad (3)$$

for $P(\xi, \tau) = P$, $\xi > 0$, $\tau \in (0, T]$.

* Corresponding author e-mail: soedeki@yahoo.com

Solution of PDEs such as (3) can be considered by semi-analytical, numerical, and approximate methods [17-20]. Goard [21] considered bond-pricing model w.r.t. group invariant solutions using classical Lie method approach. In computing the zero-coupon bonds, Sinkala *et al.* [22] applied symmetry analysis for the Vasicek and Cox-Ingersoll-Roll (CIR) models. Poee *et al.* [23] transformed the one-factor bond pricing model to one-dimensional heat equation in order to obtain fundamental solutions of zero-coupon bond models. Recently, Khalique and Motsepa [24] analysed the one-factor term structure model; thereafter, they constructed new group invariant solutions to the corresponding equation.

In this paper, the extension of (3) to time-fractional order is considered. This takes the form:

$$\begin{cases} \frac{\partial^\alpha P}{\partial \tau^\alpha} = \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P}{\partial \xi} - \xi P, \\ P(\xi, 0) = h(\xi), \alpha \in (0, 1]. \end{cases} \quad (4)$$

The fractional derivative in (4) is defined in conformable sense. For related researches on fractional, integer, financial models and solution methods, we make reference to [25-35].

In terms of organization in the remaining parts of the paper, we have in section 2 a brief note on the basic notions of conformable differential operators, section 3 is on the proposed solution method (CADM). In section 4, the proposed method is applied. Thereafter, concluding remark is presented in section 5.

2 Basic Notions of Conformable Differential Operators

For a function $\psi : [0, \infty) \rightarrow \mathbb{R}$, the conformable derivative of ψ of order $\alpha \in [0, \infty)$ is defined as:

$$C_\alpha(\psi)(t) = \lim_{\varepsilon \rightarrow 0} \left(\frac{\psi(t + \varepsilon t^{1-\alpha}) - \psi(t)}{\varepsilon} \right), \forall t \geq 0. \quad (5)$$

Reference is made to [25-28, 31] for details regarding conformable differential operators.

2.1 Properties of Conformable Differential Operators CDOs

Suppose $\psi = \psi(t)$, $\psi_1 = \psi_1(t)$ and $\psi_2 = \psi_2(t)$ are α -differentiable functions at $t > 0$, then the following holds:

$$(C1): C_\alpha(\lambda_1 \psi_1 \pm \lambda_2 \psi_2) = \left\{ \begin{matrix} \lambda_1 C_\alpha(\psi_1) \\ \pm \lambda_2 C_\alpha(\psi_2), \lambda_1, \lambda_2 \in \mathbb{R} \end{matrix} \right\}.$$

$$(C2): C_\alpha(\psi = \lambda) = 0, \lambda \in \mathbb{R}.$$

$$(C3): C_\alpha(\psi_1 \psi_2) = \psi_1 C_\alpha(\psi_2) + \psi_2 C_\alpha(\psi_1).$$

$$(C4): C_\alpha\left(\frac{\psi_1}{\psi_2}\right) = \frac{\psi_2 C_\alpha(\psi_1) - \psi_1 C_\alpha(\psi_2)}{\psi_2^2}.$$

$$(C5): C_\alpha(t^\gamma) = \gamma t^{\gamma-\alpha}, \forall \gamma \in \mathbb{R}.$$

$$(C6): C_\alpha(\psi(t)) = t^{\gamma-1} \psi'(t), \psi'(t) = \frac{d\psi(\cdot)}{dt}.$$

(C7): Suppose further that $\psi(t)$ is an n -times differentiable function at t , then:

$$C_\alpha(\psi(t)) = \begin{cases} t^{[\alpha]-\alpha} \psi^{([\alpha])}(t) \\ \alpha \in (n, n+1], \end{cases}$$

where $[\alpha]$ denotes the smallest integer such that $[\alpha] \geq \alpha$.

3 The Conformable Sense of the Decomposition Method

Let us consider a general nonlinear fractional partial differential equation (NLFDE) of the form [25-28]:

$$L_\alpha(\psi(x, t)) + R(\psi(x, t)) + N(\psi(x, t)) = \eta(x, t) \quad (6)$$

where $L_\alpha(\cdot)$ represents a linear operator based on conformable derivative of order α , with respect to t , such that $\alpha \in (n, n+1]$, R is the remaining part of the linear conformable differential operator, N denotes the nonlinear operator, while $\eta(x, t)$ is the associated non-homogeneous part (source term).

Suppose $L_\alpha(\cdot) = C_\alpha(\cdot)$ is invertible such that $L_\alpha^{-1}(\cdot)$ exists, then (6) becomes:

$$C_\alpha(\psi(x, t)) + R(\psi(x, t)) + N(\psi(x, t)) = \eta(x, t). \quad (7)$$

Hence, by the differential property of the conformable derivative (C7), we have:

$$t^{[\alpha]-\alpha} \psi^{([\alpha])}(t) + R(\psi(x, t)) + N(\psi(x, t)) = \eta(x, t). \quad (8)$$

$$\therefore t^{[\alpha]-\alpha} \frac{\partial^{[\alpha]} \psi(x, t)}{\partial t^\alpha} = \eta(x, t) - \left(\begin{matrix} R(\psi(x, t)) \\ + N(\psi(x, t)) \end{matrix} \right). \quad (9)$$

The inverse operator is defined as follows:

$$L_\alpha^{-1}(\cdot) = \int_0^t \int_0^{\tau_1} \cdots \int_0^{\tau_{n-1}} \frac{1}{\tau^{[\alpha]-\alpha}}(\cdot) d\tau_n d\tau_{n-1} \cdots d\tau_1. \quad (10)$$

So, applying (10) to both sides of (9) gives:

$$L_\alpha^{-1} \left(t^{[\alpha]-\alpha} \frac{\partial^{[\alpha]} \psi(x, t)}{\partial t^\alpha} \right) = L_\alpha^{-1} \left(\begin{matrix} \eta(x, t) \\ - \left(\begin{matrix} R(\psi(x, t)) \\ + N(\psi(x, t)) \end{matrix} \right) \end{matrix} \right).$$

$$\Rightarrow \psi(x, t) = \left\{ \begin{matrix} \psi(x, 0) + L_\alpha^{-1}(\eta(x, t)) \\ - L_\alpha^{-1}(R(\psi(x, t)) + N(\psi(x, t))) \end{matrix} \right\}. \quad (11)$$

For the decomposition of the solution, we write:

$$\psi(x, t) = \sum_{n=0}^{\infty} \psi_n(x, t) \quad (12)$$

while the nonlinear term with the Adomian polynomials A_n is defined as:

$$N(\psi(x, t)) = \sum_{n=0}^{\infty} A_n. \quad (13)$$

Note that A_n (the Adomian polynomials) is given as:

$$A_n = \frac{1}{n!} \frac{\partial^n}{\partial \xi^n} \left[N \left(\sum_{i=0}^n \xi^i h_i \right) \right]_{\xi=0}. \quad (14)$$

Thus, using (12-14) in (11) gives:

$$\sum_{n=0}^{\infty} \psi_n(x, t) = \left\{ \begin{array}{l} \psi(x, 0) + L_t^{-1} \{ \eta(x, t) \} \\ -L_t^{-1} \left\{ \left(R \left(\sum_{n=0}^{\infty} \psi_n(x, t) \right) \right) \right. \\ \left. + N \left(\sum_{n=0}^{\infty} A_n \right) \right\} \end{array} \right\}. \quad (15)$$

Hence, in recursive relation, we have:

$$\left\{ \begin{array}{l} \psi_0 = \psi(x, 0) + L_t^{-1} \{ \eta(x, t) \} \\ \psi_{n+1} = -L_t^{-1} \{ (R(\psi_n) + A_n) \}, n \geq 0. \end{array} \right. \quad (16)$$

The solution, $\psi(x, t)$ is therefore confirmed as:

$$\psi(x, t) = \lim_{n \rightarrow \infty} \sum_{n=0}^{\infty} \psi_n \quad (17)$$

4 Applications and Illustrative Examples

Here, the CADM as proposed above is applied to a Time-Fractional One-Factor Markovian Model (TF1FMM) for bond pricing as follows [3]:

$$\left\{ \begin{array}{l} \frac{\partial^\alpha P}{\partial \tau^\alpha} = \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P}{\partial \xi} - \xi P, \\ P(\xi, 0) = 1, \alpha \in (0, 1]. \end{array} \right. \quad (18)$$

Procedure: Let $D_t^\alpha P(\cdot) = C_\alpha P(\cdot)$ be applied to (18). Thus,

$$C_\alpha P(\xi, \tau) = \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P}{\partial \xi} - \xi P. \quad (19)$$

By the property: (C6), we have:

$$t^{1-\alpha} \frac{\partial P(\xi, \tau)}{\partial t} = \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P}{\partial \xi} - \xi P. \quad (20)$$

So, operating $L_\alpha^{-1}(\cdot) = \int_0^t \frac{1}{\tau^{1-\alpha}}(\cdot) d\tau$ on both sides of (20) gives:

$$P(\xi, \tau) = P(\xi, 0) + L_\alpha^{-1} \left(\frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P}{\partial \xi} - \xi P \right). \quad (21)$$

By decomposing $P(x, t)$, we have:

$$\sum_{n=0}^{\infty} P_n(\xi, \tau) = \left\{ \begin{array}{l} P(\xi, 0) \\ + L_\alpha^{-1} \left\{ \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2}{\partial \xi^2} \left(\sum_{n=0}^{\infty} P_n \right) - \xi \sum_{n=0}^{\infty} P_n \right. \\ \left. + (\beta_1 + \beta_2 \xi) \frac{\partial}{\partial \xi} \left(\sum_{n=0}^{\infty} P_n \right) \right\} \end{array} \right\}. \quad (22)$$

Thus,

$$\left\{ \begin{array}{l} P_0 = P(\xi, 0) \\ P_{n+1} = L_\alpha^{-1} \left\{ \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2}{\partial \xi^2} P_n - \xi P_n \right. \\ \left. + (\beta_1 + \beta_2 \xi) \frac{\partial}{\partial \xi} P_n \right\}, n \geq 0. \end{array} \right. \quad (23)$$


As a result, the recursive relation in (23) yields:

$$\left\{ \begin{array}{l} P_0 = P(\xi, 0) \\ P_1 = L_\alpha^{-1} \left\{ \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P_0}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P_0}{\partial \xi} - \xi P_0 \right\} \\ P_2 = L_\alpha^{-1} \left\{ \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P_1}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P_1}{\partial \xi} - \xi P_1 \right\} \\ P_3 = L_\alpha^{-1} \left\{ \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P_2}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P_2}{\partial \xi} - \xi P_2 \right\} \\ P_4 = L_\alpha^{-1} \left\{ \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P_3}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P_3}{\partial \xi} - \xi P_3 \right\} \\ \vdots \\ P_k = L_\alpha^{-1} \left\{ \frac{1}{2} \sigma^2 \xi^{2\phi} \frac{\partial^2 P_{k-1}}{\partial \xi^2} + (\beta_1 + \beta_2 \xi) \frac{\partial P_{k-1}}{\partial \xi} - \xi P_{k-1} \right\}, k \in \mathbb{N}. \end{array} \right. \quad (24)$$


Whence, for: $P(\xi, 0) = 1$, the following are obtained:

$$\left\{ \begin{array}{l} P_0 = 1, \\ P_1 = -\frac{\tau^\alpha \xi}{\alpha}, \\ P_2 = -(\beta_2 \xi - \xi^2 + \beta_1) \frac{\tau^{2\alpha}}{2! \alpha^2}, \\ P_3 = -(\beta_2^2 \xi - 3\beta_2 \xi^2 - \xi \sigma^2 + \xi^3 + \beta_1 \beta_2 - 3\beta_1 \xi) \frac{\tau^{3\alpha}}{3! \alpha^3}, \\ P_4 = \left(\begin{array}{l} -\beta_2^3 \xi + 7\beta_2^2 \xi^2 + 4\xi \sigma^2 \beta_2 \\ -6\beta_2 \xi^3 - 4\xi^2 \sigma^2 + \xi^4 - \beta_1 \beta_2^2 \\ +10\beta_1 \beta_2 \xi + \beta_1 \sigma^2 - 6\beta_1 \xi^2 + 3\beta_1^2 \end{array} \right) \frac{\tau^{4\alpha}}{4! \alpha^4}, \\ P_5 = \left(\begin{array}{l} -\beta_2^4 \xi + 15\beta_2^3 \xi^2 + 11\beta_2^2 \sigma^2 \xi \\ -25\beta_2^2 \xi^3 - 30\beta_2 \sigma^2 \xi^2 + 10\beta_2 \xi^4 \\ -4\sigma^4 \xi + 10\sigma^2 \xi^3 - \xi^5 - \beta_1 \beta_2^3 \\ +25\beta_1 \beta_2^2 \xi + 4\beta_1 \beta_2 \sigma^2 \\ -40\beta_1 \beta_2 \xi^2 - 15\beta_1 \sigma^2 \xi + 10\beta_1 \xi^3 \\ +10\beta_1^2 \beta_2 - 15\beta_1^2 \xi \end{array} \right) \frac{\tau^{5\alpha}}{5! \alpha^5}, \\ P_6 = - \left(\begin{array}{l} \beta_2^5 \xi - 31\beta_2^4 \xi^2 - 26\beta_2^3 \sigma^2 \xi \\ +90\beta_2^3 \xi^3 + 146\beta_2^2 \sigma^2 \xi^2 \\ -65\beta_2^2 \xi^4 + 34\beta_2 \sigma^4 \xi - 120\beta_2 \sigma^2 \xi^3 \\ +15\beta_2 \xi^5 - 34\sigma^4 \xi^2 + 20\sigma^2 \xi^4 - \xi^6 \\ +\beta_1 \beta_2^4 - 56\beta_1 \beta_2^3 \xi - 11\beta_1 \beta_2^2 \sigma^2 \\ +180\beta_1 \beta_2^2 \xi^2 + 119\beta_1 \beta_2 \sigma^2 \xi \\ -110\beta_1 \beta_2 \xi^3 + 4\beta_1 \sigma^4 - 75\beta_1 \sigma^2 \xi^2 \\ +15\beta_1 \xi^4 - 25\beta_1^2 \beta_2^2 + 105\beta_1^2 \beta_2 \xi \\ +15\beta_1^2 \sigma^2 - 45\beta_1^2 \xi^2 + 15\beta_1^3 \end{array} \right) \frac{\tau^{6\alpha}}{6! \alpha^6}, \end{array} \right.$$

$$P_7 = - \left(\begin{array}{l} \beta_2^6 \xi - 63\beta_2^5 \xi^2 - 57\beta_2^4 \sigma^2 \xi + 301\beta_2^4 \xi^3 \\ + 588\beta_2^3 \sigma^2 \xi^2 - 350\beta_2^3 \xi^4 + 180\beta_2^2 \sigma^4 \xi \\ - 896\beta_2^2 \sigma^2 \xi^3 + 140\beta_2^2 \xi^5 - 462\beta_2 \sigma^4 \xi^2 \\ + 350\beta_2 \sigma^2 \xi^4 - 21\beta_2 \xi^6 - 34\sigma^6 \xi \\ + 154\sigma^4 \xi^3 - 35\sigma^2 \xi^5 + \xi^7 + \beta_1 \beta_2^5 \\ - 119\beta_1 \beta_2^4 \xi - 26\beta_1 \beta_2^3 \sigma^2 + 686\beta_1 \beta_2^3 \xi^2 \\ + 602\beta_1 \beta_2^2 \sigma^2 \xi - 770\beta_1 \beta_2^2 \xi^3 \\ + 34\beta_1 \beta_2 \sigma^4 - 959\beta_1 \beta_2 \sigma^2 \xi^2 + 245\beta_1 \beta_2 \xi^4 \\ - 147\beta_1 \sigma^4 \xi + 245\beta_1 \sigma^2 \xi^3 - 21\beta_1 \xi^5 \\ - 56\beta_1^2 \beta_2^3 + 490\beta_1^2 \beta_2^2 \xi + 119\beta_1^2 \beta_2 \sigma^2 \\ - 525\beta_1^2 \beta_2 \xi^2 - 210\beta_1^2 \sigma^2 \xi + 105\beta_1^2 \xi^3 \\ + 105\beta_1^3 \beta_2 - 105\beta_1^3 \xi \end{array} \right) \frac{\tau^7 \alpha}{7! \alpha^7},$$

Fig. 1: Approximate analytical solution graphic for $\alpha = 1$

$$P_8 = \left(\begin{array}{l} -\beta_2^7 \xi + 127\beta_2^6 \xi^2 + 120\beta_2^5 \sigma^2 \xi - 966\beta_2^5 \xi^3 \\ - 2136\beta_2^4 \sigma^2 \xi^2 + 1701\beta_2^4 \xi^4 - 768\beta_2^3 \sigma^4 \xi \\ + 5376\beta_2^3 \sigma^2 \xi^3 - 1050\beta_2^3 \xi^5 + 3792\beta_2^2 \sigma^4 \xi^2 \\ - 3696\beta_2^2 \sigma^2 \xi^4 + 266\beta_2^2 \xi^6 + 496\beta_2 \sigma^6 \xi \\ - 3024\beta_2 \sigma^4 \xi^3 + 840\beta_2 \sigma^2 \xi^5 - 28\beta_2 \xi^7 \\ - 496\sigma^6 \xi^2 + 504\sigma^4 \xi^4 - 56\sigma^2 \xi^6 + \xi^8 - \beta_1 \beta_2^6 \\ + 246\beta_1 \beta_2^5 \xi + 57\beta_1 \beta_2^4 \sigma^2 - 2394\beta_1 \beta_2^4 \xi^2 \\ - 2490\beta_1 \beta_2^3 \sigma^2 \xi + 4396\beta_1 \beta_2^3 \xi^3 - 180\beta_1 \beta_2^2 \sigma^4 \\ + 7518\beta_1 \beta_2^2 \sigma^2 \xi^2 - 2450\beta_1 \beta_2^2 \xi^4 \\ + 2064\beta_1 \beta_2 \sigma^4 \xi - 4564\beta_1 \beta_2 \sigma^2 \xi^3 + 476\beta_1 \beta_2 \xi^5 \\ + 34\beta_1 \sigma^6 - 1344\beta_1 \sigma^4 \xi^2 + 630\beta_1 \sigma^2 \xi^4 - 28\beta_1 \xi^6 \\ + 119\beta_1^2 \beta_2^4 - 1918\beta_1^2 \beta_2^3 \xi - 602\beta_1^2 \beta_2^2 \sigma^2 \\ + 3850\beta_1^2 \beta_2^2 \xi^2 + 2772\beta_1^2 \beta_2 \sigma^2 \xi - 1820\beta_1^2 \beta_2 \xi^3 \\ + 147\beta_1^2 \sigma^4 - 1260\beta_1^2 \sigma^2 \xi^2 + 210\beta_1^2 \xi^4 \\ - 490\beta_1^3 \beta_2^2 + 1260\beta_1^3 \beta_2 \xi + 210\beta_1^3 \sigma^2 \\ - 420\beta_1^3 \xi^2 + 105\beta_1^4 \end{array} \right) \kappa,$$

Fig. 2: Approximate analytical solution graphic for $\alpha = 0.5$


where $\kappa = \frac{\tau^8 \alpha}{8! \alpha^8}$.
Therefore,

$$P(\xi, \tau) = \lim_{k \rightarrow \infty} P_k = \sum_{n=0}^{\infty} P_k. \quad (25)$$

Equation (25) is an approximate-analytical solution of (18) corresponding to the time-fractional one-factor Markovian model for bond pricing. We consider the financial data according to [1] w.r.t CIR model with $\sigma = 0.0894$, $\beta_1 = 0.00315$, and $\beta_2 = -0.0555$. The plots of the resulting solutions for $\alpha = 1$, $\alpha = 0.5$, and $\alpha = 0.75$ are displayed in Fig.1, Fig. 2, and Fig. 3 respectively.

4.1 Numerical Solutions

This subsection deals significantly with the approximate-analytical solutions obtained via the proposed method (CADM). In the classical setting when

Fig. 3: Approximate analytical solution graphic for $\alpha = 0.75$

the parameter value, $\alpha \rightarrow 1$, Fig. 1 describes the geometrical behaviour of the approximate solution in line with the market (financial) data as contained in [1]. In a similar way, Fig. 2 and Fig.3 describe the geometrical behaviour of the approximate solution for the time-orders, $\alpha \rightarrow 0.5$ and $\alpha \rightarrow 0.75$ respectively. These time fractional cases ($\alpha \rightarrow 0.5$ and $\alpha \rightarrow 0.75$) are compared with the integer case ($\alpha \rightarrow 1$) making the result in [1] a particular case of this present work.

5 Prospectives

In this paper, approximate-analytical solution is proposed for a Time-Fractional One-Factor Markovian Model (TF1FMM) for bond pricing whose derivative is defined in conformable sense. Preference is given to the CIR model version of TF1FMM with the market parameters according to [1] in order to clarify the effectiveness of the proposed solution method. Thereafter, the solutions at different values of time-fractional order are graphically displayed. This proposed method of solution can be extended to multi-factor models formulated in terms of stochastic dynamics. In addition, the solution obtained, and the proposed method can effectively serve as benchmarks for further researches in related areas via other semi-analytical methods.

Conflict of Interests

The authors declare no conflict of interest regarding this paper.

Acknowledgement

The authors¹ express sincere thanks to Covenant University for all forms of support including the provision of good working environment. Also, the constructive comments of the anonymous referees are highly appreciated.

References

- [1] B. Stehlikova, A simple analytic approximation formula for the bond price in the Chan-Karolyi-Longstaff-Sanders model, *International Journal of Numerical Analysis and Modeling*, Series B, **4**, 224-234, (2013).
- [2] Y. Ait-Sahalia, Testing continuous-time models of the spot interest rate, *Review of Financial Studies*, **9**, 385-426, (1996).
- [3] Y. Ait-Sahalia, Transition densities for interest rate and other nonlinear diffusions, *Journal of Finance*, **54**, 1361-1395, (1999).
- [4] F. Black, M. Scholes, The pricing of options and corporate liabilities, *Journal of Political Economy*, **81**, 637-654, (1973).
- [5] Z. Buckova, B. Stehlikova, D. Sevcovic, Numerical and analytical methods for bond pricing in short rate convergence models of interest rates, <https://arxiv.org/pdf/1607.04968.pdf>, (2016).
- [6] S. Shreve, *Stochastic Calculus for Finance II: Continuous Time Models*, Springer, New York, (2004).
- [7] K.L. Chan, G.A. Karolyi, F.A. Longstaff, and A.B. Sanders, An empirical comparison of alternative models of the short-term interest rate, *Journal of Finance*, **47**, 1209-1227, (1992).
- [8] J.C. Cox, J.E. Ingersoll, S. A. Ross, An analysis of variable rate loan contracts, *Journal of Finance*, **35**, 389-403, (1980).
- [9] R. C. Merton, Theory of rational option pricing, *The Bell Journal of Economics and Management Science*, **4**, 141-183, (1973).
- [10] O.A. Vasicek, An equilibrium characterization of the term structure, *Journal of Financial Economics*, **5**, 177-188, (1977).
- [11] M.J. Brennan, Michael, E.S. Schwartz, Savings bonds, retractable bonds, and callable bonds, *Journal of Financial Economics*, **3**, 133-155, (1977).
- [12] U.L. Dothan, On the term structure of interest rates, *Journal of Financial Economics*, **6**, 59-69, (1978).
- [13] M.J. Brennan, Michael, E.S. Schwartz, Analyzing convertible bonds, *Journal of Financial and Quantitative Analysis*, **15**, 907-929, (1980).
- [14] G. Courtadon, The pricing options on default-free bonds, *Journal of Financial and Quantitative Analysis*, **17**, 75-100, (1982).
- [15] J.C. Cox, J.E., Ingersoll, S.A. Ross, A theory of the term structure of interest rates, *Econometrica*, **53**, 385-408, (1985).
- [16] S.O. Edeki, O.O. Ugbebor, and E. A. Owoloko, On a dividend-paying stock options pricing model (SOPM) using constant elasticity of variance stochastic dynamics, *International Journal of Pure and Applied Mathematics*, **106**, 1029-1036, (2016).
- [17] G.O. Akinlabi, S.O. Edeki, On approximate and closed-form solution method for initial-value wave-like models, *International Journal of Pure and Applied Mathematics*, **107**, 449-456 (2016).
- [18] J. Biazar and F. Goldoust, The Adomian decomposition method for the Black-Scholes equation, 3rd *International Conference on Applied Mathematics and Pharmaceutical Sciences* (ICAMP 2013), 321-323, (2013).
- [19] J.G. Oghonyon, T.O. Mosaku, M. RemilekunOdekunle, O.F. Imaga, T. Abodunrin, Mathematica computer programming codes of exponentially fitted concurrent Milne's device for solving special problems, *International Journal of Mechanical Engineering and Technology*, **9**, 1577-1592, (2018).
- [20] A.A. Ayodele and K.A. Charles, Portfolio selection problem using generalized differential evolution 3, *Applied Mathematical Sciences*, **9**, 2069-2082, (2015).
- [21] J. Goard, New solutions to the bond-pricing equation via Lie's classical method, *Math. Comput. Modelling*, **32**, 299-313, (2000).
- [22] W. Sinkala, P.G.L. Leach, J.G. O'Hara, Zero-coupon bond prices in the Vasicek and CIR models: Their computation as group-invariant solutions, *Math Method Appl. Sci.* **31**, 665-678, (2008).
- [23] C.A. Poore, F.M. Mahomed, C.W. Soh, Fundamental solutions for zero-coupon bond pricing models, *Nonlinear Dynam.* **36**, 69-76, (2004).

- [24] C.M. Khalique, T. Motsepa, Lie symmetries, group-invariant solutions and conservation laws of the Vasicek pricing equation of mathematical finance, *Physica A: Statistical Mechanics and its Applications*, **505**, 871-879, (2018).
- [25] R. Khalil, M. Al Horani, A. Yousef, M. Sababheh, A new definition of fractional derivative, *J. Comput. Appl. Math.*, **264**, 65-70, (2014).
- [26] T. Abdeljawad, On conformable fractional calculus, *J. Comput. Appl. Math.*, **279**, 57-66 (2015).
- [27] O. Acan and D. Baleanu, A new numerical technique for solving fractional partial differential equations, *Miskolc Mathematical Notes*, **19**, 3-18, (2018).
- [28] M. Yavuz, Novel solution methods for initial boundary value problems of fractional order with conformable differentiation, *An International Journal of Optimization and Control: Theories & Applications*, **8**, 1-7, (2018).
- [29] S.O. Edeki, O.O. Ugbebor, and E.A. Owoloko, Analytical Solution of the time-fractional order Black-Scholes model for stock option valuation on no dividend yield basis, *IAENG International Journal of Applied Mathematics*, **47** (4), 407-416, (2017).
- [30] S.O. Edeki, E. A. Owoloko, and O.O. Ugbebor, The modified Black-Scholes model via constant elasticity of variance for stock options valuation, *AIP Conference Proceedings*, **1705**, 4940289, (2016).
- [31] S.O. Edeki, G.O. Akinlabi, Local fractional operator for a one-dimensional coupled burger equation of non-integer time order parameter, *Journal of Mathematical and Fundamental Sciences*, **50**, 28-39, (2018).
- [32] K. Shah, S. Zeb, R.A. Khan, Multiplicity results of multi-point boundary value problem of nonlinear fractional differential equations, *Applied Mathematics and Information Sciences*, **12**, 727-734, (2018).
- [33] S.O. Edeki, O.O. Ugbebor, and E. A. Owoloko, He's polynomials for analytical solutions of the Black-Scholes pricing model for stock option valuation, *Lecture Notes in Engineering and Computer Science*, **2224**, 632-634, (2016).
- [34] E. Ozbilge, A. Demir, An inverse source problem in time-space fractional differential equations, *Applied Mathematics and Information Sciences*, **12**, 587-591, (2018).
- [35] R.M. Jena, S. Chakraverty, A new iterative method based solution for fractional Black-Scholes option pricing Equations (BSOPE), *SN Appl. Sci.*, **1**: 95, (2019).



S. O. Edeki currently works as a Lecturer and Researcher at the Department of Mathematics, Covenant University, Ota, Nigeria. His area of interest includes Stochastic Analysis and its applications to finance (Financial Mathematics), Fractional Calculus, Probability theory, Numerical Solutions of (Stochastic) Differential Equations, and Dynamical Systems. His current project is on 'Improved Black-Scholes Option Pricing Model'. He has published good number of journal and conference articles in well-indexed outlets of high international standard abstracted in Scopus and Web of Science.



I. Adinya is a Lecturer at the Department of Mathematics, University of Ibadan, Nigeria. Her areas of interest are in Financial and Computational Mathematics. She is a member of Nigerian Mathematical Society (NMS) and Society of Industrial and Applied Mathematics (SIAM). She has published research articles in reputed international journals of mathematical sciences.



G. O. Akinlabi is a Lecturer and Researcher at the Department of Mathematics, Covenant University, Ota, Nigeria. She obtained both her first and second degrees (B.Sc. & M.Sc.) in Mathematics from University of Ilorin, Ilorin, Nigeria. She is currently pursuing her Ph.D (Mathematics) at Covenant University. Her area of interest includes Numerical Analysis, Differential Equations and Complex Analysis. She has published reasonable number of papers in well-indexed international journals.



O. P. Ogundile received his Master's Degree (M.Sc Mathematics) from the prestigious University of Ibadan, Nigeria and his B.Sc from Covenant University, Nigeria. Currently, he is a faculty member at Covenant University. His area of interest includes Uncertain Differential Equation, and Financial Mathematics. He has published research articles in reputed international journals of mathematical sciences.