

1-1-2021

A third-order p-laplacian boundary value problem on an unbounded domain

SAMUEL AZUBUIKE IYASE

OGBU FAMOUS IMAGA

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

IYASE, SAMUEL AZUBUIKE and IMAGA, OGBU FAMOUS (2021) "A third-order p-laplacian boundary value problem on an unbounded domain," *Turkish Journal of Mathematics*: Vol. 45: No. 6, Article 2.

<https://doi.org/10.3906/mat-2002-89>

Available at: <https://dctubitak.researchcommons.org/math/vol45/iss6/2>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

A third-order p -laplacian boundary value problem on an unbounded domain

Samuel Azubuikwe IYASE[✉], Ogbu Famous IMAGA*[✉]
 Department of Mathematics, Covenant University, Ota, Ogun State, Nigeria

Received: 21.02.2020

Accepted/Published Online: 16.08.2021

Final Version: 29.11.2021

Abstract: In this work, we apply Leray-Schauder continuation principle to establish the existence of at least one solution to the third order p -Laplacian boundary value problem on an unbounded domain of the form

$$(w(t)\varphi_p(u''(t)))' = K(t, u(t), u'(t), u''(t)), t \in (0, \infty)$$

$$u(0) = 0, u'(0) = \sum_{i=1}^m \alpha_i \int_0^{\xi_i} u(t)dt, \lim_{t \rightarrow \infty} (w(t)\varphi_p(u''(t))) = 0$$

under the nonresonant condition $\sum_{i=1}^m \alpha_i \xi_i^2 \neq 2$.

Key words: Nonresonance, p -Laplacian, unbounded domain, third-order, boundary value problem

1. Introduction

The purpose of this paper is to obtain existence of at least one solution for the third order p -Laplacian boundary value problem

$$(w(t)\varphi_p(u''(t)))' = K(t, u(t), u'(t), u''(t)), t \in (0, \infty) \quad (1.1)$$

$$u'(0) = \sum_{i=1}^m \alpha_i \int_0^{\xi_i} u(t)dt, u(0) = 0, \lim_{t \rightarrow \infty} (w(t)\varphi_p(u''(t))) = 0 \quad (1.2)$$

where

$\varphi_p(s) = |s|^{p-2}s$, $K : [0, \infty) \times \mathfrak{R}^3 \rightarrow \mathfrak{R}$ is a Caratheodory function with respect to $L^1[0, \infty)$, $\alpha_i \in \mathfrak{R} (1 \leq i \leq m)$, $0 < \xi_1 < \xi_2 < \dots < \xi_m < 1$. $w(t) > 0, t \in [0, \infty)$, $w \in C[0, \infty) \cap C^1[0, \infty)$, $\frac{1}{w} \in L^1[0, \infty)$ and $\sum_{i=1}^m \alpha_i \xi_i^2 \neq 2$.

The condition $\sum_{i=1}^m \alpha_i \xi_i^2 \neq 2$ is critical since we require a trivial kernel for the differential operator. The boundary value problem (1.1)–(1.2) is then said to be at nonresonance. In [6] we proved the existence of solutions for the boundary value problem.

$$(q(t)u''(t))' = f(t, u(t), u'(t), u''(t)), t \in (0, \infty) \quad (1.3)$$

$$u'(0) = \sum_{i=1}^m \alpha_i \int_0^{\xi_i} u(t)dt, u(0) = 0, \lim_{t \rightarrow \infty} q(t)u''(t) = 0 \quad (1.4)$$

*Correspondence: imaga.ogbu@covenantuniversity.edu.ng

2010 AMS Mathematics Subject Classification: 34B10, 34B15

under the resonant condition $\sum_{i=1}^m \alpha_i \xi_i^2 = 2$.

The method of investigation in [6] was based on coincidence degree arguments. In this work, we shall utilise topological degree methods based on the Leray-Schauder degree theory [10]. The contribution of this paper is to extend the results of (1.3)–(1.4) to a p -Laplacian boundary value problem where the differential operator has trivial solutions.

Boundary value problems on an unbounded domain with integral boundary conditions have various applications. For example they are used to model, the steady flow of gas through a semi-infinite porous medium, theory of drain flows, blood flow models. Boundary value problems with p -Laplacian operators also have applications in non-Newtonian mechanics, nonlinear elasticity, glaciology, etc.

For some recent works where Leray-Schauder principle has been applied for boundary value problems on unbounded domain or half-line see [1, 2, 4, 6–8] and references therein.

This paper is organized as follows. In Section 2 we present some background definitions, theorems and a-priori estimates that will be used in the proof of the main existence results. In Section 3 we prove the main existence result. Section 4 will be devoted to providing an example to demonstrate our results.

2. Preliminaries

In what follows, we shall use the following definition and lemmas.

Definition 2.1 *The mapping $f : [0, \infty) \times \mathbb{R}^n \rightarrow \mathbb{R}$ is an $L^1[0, \infty)$ -Caratheodory if the following conditions hold*

- (i) *for each $z \in \mathbb{R}^n$, $f(t, z)$ is Lebesgue measurable;*
- (ii) *for a.e $t \in (0, \infty)$, $f(t, z)$ is a continuous on \mathbb{R}^n ;*
- (iii) *for each $r > 0$ there exists $\alpha_r \in (0, \infty)$ such that for a.e $t \in [0, \infty)$ and every z such that $|z| \leq r$ we have $|f(t, z)| \leq \alpha_r(t)$*

Let $AC[0, \infty)$ denote the space of absolutely continuous functions on $[0, \infty)$. In what follows we shall utilise the following spaces

$$X = \left\{ u \in C^2[0, \infty), (w\varphi_p(u''))' \in AC[0, \infty), \lim_{t \rightarrow \infty} e^{-t} |u^{(i)}(t)| \text{ exists for } 0 \leq i \leq 2, (w\varphi_p(u''))' \in L^1[0, \infty) \right\}, \tag{2.1}$$

$$\|u\| = \max \left[\sup_{t \in [0, \infty)} e^{-t} |u(t)|, \sup_{t \in [0, \infty)} e^{-t} |u'(t)|, \sup_{t \in [0, \infty)} e^{-t} |u''(t)| \right]. \tag{2.2}$$

Thus X is a Banach Space . Let $Z = L^1[0, \infty)$ with the norm

$$\|y\|_1 = \int_0^\infty |y(t)| dt, \quad y \in Z \tag{2.3}$$

Define the mapping $N : X \rightarrow Z$ by

$$Nu(t) = K(t, u(t), u'(t), u''(t)), \quad t \in [0, \infty) \tag{2.4}$$

In this work, we shall utilise the Leray-Schauder continuation principle which has been applied in many papers in the literature. However since $[0, \infty)$ is not compact, the compactness principle on the bounded interval $[0, 1]$ cannot be applied here. We shall rely on the following compactness criterion.

Lemma 2.2 [1] *Let $H \subset X$. Then H is relatively compact in X if the following conditions are satisfied*

- (i) H is bounded in X ;
- (ii) The family $W^i = \{\psi_i : \psi_i(t) = e^{-t}u^i(t), t \geq 0, u \in H\}$ is equicontinuous on any compact subinterval of $[0, \infty)$ for $i = 0, 1, \dots, n - 1$;
- (iii) The family $W^i = \{\psi_i : \psi_i(t) = e^{-t}u^i(t), t \geq 0, u \in H\}$ is equiconvergent at infinity for $i = 0, 1, \dots, n - 1$.

We shall need the following constants

$$A = e^{-1} \left\| \frac{1}{w} \right\|_1^{p-1} + e^{-1} \left\| \frac{1}{w} \right\|_1^{p-1} \frac{1}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \quad (2.5)$$

$$B = \left\| \frac{1}{w} \right\|_1^{p-1} + \left\| \frac{1}{w} \right\|_1^{p-1} \frac{1}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \quad (2.6)$$

$$C = \left\| \frac{1}{w} \right\|_\infty^{p-1}. \quad (2.7)$$

Lemma 2.3 *Let $h \in Z$. Then the unique solution of the equation.*

$$(w(t)\varphi_p(u''))' = h(t), t \in [0, \infty) \quad (2.8)$$

subject to (1.2) is

$$u(t) = - \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds + \frac{-2t}{2 - \sum_{i=1}^m \alpha_i \xi_i^2} \sum_{i=1}^m \alpha_i \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds dt. \quad (2.9)$$

Proof From (2.8) and (1.2) we derive

$$\varphi_p(u''(t)) = - \frac{1}{w(t)} \int_t^\infty h(\tau) d\tau,$$

$$u'' = -\varphi_q \left(\frac{1}{w(t)} \right) \varphi_q \int_t^\infty h(\tau) d\tau$$

and hence

$$u(t) = - \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds + tu'(0).$$

Therefore from (1.2) we obtain

$$u'(0) = \frac{-2 \sum_{i=1}^m \alpha_i}{2 - \sum_{i=1}^m \alpha_i \xi_i^2} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds dt.$$

Hence (2.9) follows. □

Lemma 2.4 *Let $h \in Z$. Then the solution (2.9) satisfies*

$$e^{-t} |u(t)| \leq A \|h\|_1^{\frac{1}{p-1}}, \tag{2.10}$$

$$e^{-t} |u'(t)| \leq B \|h\|_1^{\frac{1}{p-1}} \tag{2.11}$$

and

$$e^{-t} |u''(t)| \leq C \|h\|_1^{\frac{1}{p-1}}. \tag{2.12}$$

Hence,

$$\|u\| \leq \max(A, B, C) \|h\|_1^{\frac{1}{p-1}}. \tag{2.13}$$

Proof From (2.9) we obtain

$$\begin{aligned} e^{-t} |u(t)| &= e^{-t} \left| - \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) \right. \\ &\quad \left. \frac{-2t \sum_{i=1}^m \alpha_i}{2 - \sum_{i=1}^m \alpha_i \xi_i^2} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds dt \right| \\ &\leq \sup_{t \in [0, \infty)} e^{-t} \left[\int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |h(\tau)| d\tau \right) dr ds \right. \\ &\quad \left. + \frac{2t \sum_{i=1}^m |\alpha_i|}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |h(\tau)| d\tau \right) dr ds dt \right] \\ &\leq \left[\sup_{t \in [0, \infty)} e^{-t} t \left\| \frac{1}{w} \right\|_1^{\frac{1}{p-1}} \right. \\ &\quad \left. + 2 \sup_{t \in [0, \infty)} e^{-t} t \frac{\sum_{i=1}^m |\alpha_i|}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) dr ds dt \right] \|h\|_1^{\frac{1}{p-1}} \\ &= \left[e^{-1} \left\| \frac{1}{w} \right\|_1^{\frac{1}{p-1}} + e^{-1} \left\| \frac{1}{w} \right\|_1^{\frac{1}{p-1}} \frac{\sum_{i=1}^m |\alpha_i| \xi_i^2}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \right] \|h\|_1^{\frac{1}{p-1}}, \end{aligned}$$

using (2.5), we get

$$\begin{aligned}
 e^{-t} |u(t)| &= A \|h\|_1^{\frac{1}{p-1}} \tag{2.14} \\
 e^{-t} |u'(t)| &= e^{-t} \left| - \int_0^t \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr, \right. \\
 &\quad \left. - \frac{2 \sum_{i=1}^m \alpha_i}{2 - \sum_{i=1}^m \alpha_i \xi_i^2} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds dt \right| \\
 &\leq \sup_{t \in (0, \infty)} e^{-t} \left[\int_0^t \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |h(\tau)| \right) d\tau dr \right. \\
 &\quad \left. + \frac{2 \sum_{i=1}^m |\alpha_i|}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |h(\tau) d\tau| \right) dr ds dt \right] \\
 &\leq \left[\left\| \frac{1}{w} \right\|_1^{\frac{1}{p-1}} + \left\| \frac{1}{w} \right\|_1^{\frac{1}{p-1}} \frac{\sum_{i=1}^m |\alpha_i| \xi_i^2}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \right] \|h\|_1^{\frac{1}{p-1}},
 \end{aligned}$$

and from (2.6), we have

$$\begin{aligned}
 e^{-t} |u'(t)| &= B \|h\|_1^{\frac{1}{p-1}}, \tag{2.15} \\
 e^{-t} |u''(t)| &= e^{-t} \left| \varphi_q \left(\frac{1}{w(t)} \right) \varphi_q \left(\int_t^\infty h(\tau) d\tau \right) \right| \\
 &\leq \sup_{t \in [0, \infty)} e^{-t} \left| \varphi_q \left(\frac{1}{|w(t)|} \right) \varphi_q \left(\int_t^\infty |h(\tau) d\tau| \right) \right| \\
 &\leq \left\| \frac{1}{w} \right\|_\infty^{\frac{1}{p-1}} \|h\|_1^{\frac{1}{p-1}},
 \end{aligned}$$

from (2.7), we have

$$e^{-t} |u''(t)| = C \|h\|_1^{\frac{1}{p-1}}. \tag{2.16}$$

Hence

$$\|u\| \leq \max(A, B, C) \|h\|_1^{\frac{1}{p-1}} = D \|h\|_1^{\frac{1}{p-1}}. \tag{2.17}$$

□

Define $M : Z \rightarrow X$ by

$$\begin{aligned}
 (Mh)(t) &= - \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds \\
 &\quad - \frac{2t \sum_{i=1}^m \alpha_i}{2 - \sum_{i=1}^m \alpha_i \xi_i^2} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds dt.
 \end{aligned}$$

Then M is well defined for all $h \in Z$ and is the unique solution of the differential equation

$$(w(t)\varphi_p(u''(t)))' = h(t) \text{ a. e } t \in (0, \infty),$$

subject to the boundary condition (1.2).

Let $T = MN : X \rightarrow X$ be defined by

$$\begin{aligned} Tu(t) = & - \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds \\ & - \frac{2t \sum_{i=1}^m \alpha_i}{2 - \sum_{i=1}^m \alpha_i \xi_i^2} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty h(\tau) d\tau \right) dr ds dt \end{aligned}$$

for a.e $t \in [0, \infty)$. Then T is well defined. We use the compactness criterion in Lemma 2.1 to prove that T is compact.

Lemma 2.5 *The mapping $T : X \rightarrow X$ is compact.*

Proof Let $H \subset X$ be bounded i.e there exists $r > 0$ such that $|u| < r$ for all $u \in H$. Since $K : [0, \infty) \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is $L^1[0, \infty)$ Caratheodory there exists $\alpha_r \in Z$ such that $|Nu(t)| \leq \alpha_r(t)$ for all $t \in [0, \infty)$ and all $u \in H$.

For $u \in H$, we have

$$\begin{aligned} e^{-t} |Tu(t)| = & e^{-t} \left| - \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_0^\infty Nu(\tau) d\tau \right) dr ds \right. \\ & \left. - \frac{2t \sum_{i=1}^m \alpha_i}{(2 - \sum_{i=1}^m \alpha_i \xi_i^2)} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty Nu(\tau) d\tau \right) dr ds dt \right| \\ \leq & \sup_{t \in [0, \infty)} e^{-t} \left[\int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \int_r^\infty (|Nu(\tau)| d\tau) dr ds \right. \\ & \left. + \frac{2t \sum_{i=1}^m |\alpha_i|}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |Nu(\tau) d\tau \right) dr ds dt \right] \\ \leq & A \| \alpha_r \|_1^{\frac{1}{p-1}}, \end{aligned} \tag{2.18}$$

$$\begin{aligned} e^{-t} |(Tu)'(t)| = & e^{-t} \left| - \int_0^t \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty Nu(\tau) d\tau \right) dr \right. \\ & \left. - \frac{2 \sum_{i=1}^m \alpha_i}{(2 - \sum_{i=1}^m \alpha_i \xi_i^2)} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty Nu(\tau) d\tau \right) dr ds dt \right| \\ \leq & \sup_{t \in [0, \infty)} e^{-t} \left[\int_0^t \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |Nu(\tau)| d\tau \right) dr \right. \\ & \left. + \frac{2 \sum_{i=1}^m |\alpha_i|}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |Nu(\tau)| d\tau \right) dr ds dt \right] \\ \leq & B \| \alpha_r \|_1^{\frac{1}{p-1}}, \end{aligned} \tag{2.19}$$

$$\begin{aligned}
 e^{-t} \left| (Tu)''(t) \right| &= e^{-t} \left| -\varphi_q \left(\frac{1}{w(t)} \right) \varphi_q \left(\int_t^\infty Nu(\tau) d\tau \right) \right| \\
 &\leq \sup_{t \in [0, \infty)} e^{-t} \varphi_q \left(\frac{1}{|w(t)|} \right) \varphi_q \left(\int_t^\infty |Nu(\tau)| d\tau \right) \\
 &\leq C \|\alpha_r\|_1^{\frac{1}{p-1}}.
 \end{aligned} \tag{2.20}$$

(2.18), (2.19) and (2.20) implies that

$$\|Tu\| \leq \max(A, B, C) \|\alpha_r\|_1^{\frac{1}{p-1}} = D \|\alpha_r\|_1^{\frac{1}{p-1}} \tag{2.21}$$

Thus $T(H)$ is bounded in X .

To prove equicontinuity of $T(H)$, let $t_1, t_2 \in [0, L]$ where $L \in (0, \infty)$ with $t_1 < t_2$ then

$$\begin{aligned}
 \left| e^{-t_2}(Tu)(t_2) - e^{-t_1}(Tu)(t_1) \right| &= \left| \int_{t_1}^{t_2} [e^{-s}(Tu)(s)]' ds \right| \\
 &= \left| \int_{t_1}^{t_2} -e^{-s} [Tu](s) ds + \int_{t_1}^{t_2} e^{-s} [Tu]'(s) ds \right| \\
 &\leq \int_{t_1}^{t_2} e^{-s} |Tu|(s) ds + \int_{t_1}^{t_2} e^{-s} |[Tu]'(s)| ds \\
 &\leq 2(t_2 - t_1) \|Tu\| \leq 2D(t_2 - t_1) \|\alpha_r\|_1^{\frac{1}{p-1}} \rightarrow 0 \text{ as } t_1 \rightarrow t_2.
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 \left| e^{-t_2}(Tu)'(t_2) - e^{-t_1}(Tu)'(t_1) \right| &= \left| \int_{t_1}^{t_2} [e^{-s}(Tu)'(s)]' ds \right| \\
 &= \left| \int_{t_1}^{t_2} [-e^{-s}(Tu)'(s) + e^{-s}(Tu)''(s)] ds \right| \\
 &\leq 2(t_2 - t_1) \|Tu\| \leq 2(t_2 - t_1) D \|\alpha_r\|_1^{\frac{1}{p-1}} \rightarrow 0 \text{ as } t_1 \rightarrow t_2,
 \end{aligned}$$

and

$$\begin{aligned}
 \left| e^{-t_2} \varphi_p(Tu)''(t_2) - e^{-t_1} \varphi_p(Tu)''(t_1) \right| &= \left| -e^{-t_2} \frac{1}{w(t_2)} \int_{t_2}^\infty Nu(\tau) d\tau + e^{-t_1} \frac{1}{w(t_1)} \int_{t_1}^\infty Nu(\tau) d\tau \right| \\
 &= \left| \left(\frac{e^{-t_1}}{w(t_1)} - \frac{e^{-t_2}}{w(t_2)} \right) \int_{t_2}^\infty Nu(\tau) d\tau - \frac{e^{-t_1}}{w(t_1)} \int_{t_2}^\infty Nu(\tau) d\tau + \frac{e^{-t_1}}{w(t_1)} \int_{t_1}^\infty Nu(\tau) d\tau \right| \\
 &\leq \left| \frac{e^{-t_1}}{w(t_1)} - \frac{e^{-t_2}}{w(t_2)} \right| \|\alpha_r\|_1 + \frac{e^{-t_1}}{|w(t_1)|} \int_{t_1}^{t_2} |\alpha_r(\tau)| d\tau \rightarrow 0 \text{ as } t_1 \rightarrow t_2.
 \end{aligned}$$

Therefore $|e^{-t_2}(Tu)''(t_2) - e^{-t_1}(Tu)''(t_1)| \rightarrow 0$ as $t_1 \rightarrow t_2$.

The above computations shows that the set $T(H)$ is equicontinuous on every compact subinterval of $[0, \infty)$

Next we show that $T(H)$ is equiconvergent at infinity. For $u \in H$, we have

$$\begin{aligned}
 e^{-t} |(Tu)(t)| &\leq e^{-t} \left[\int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |Nu(\tau)| d\tau \right) \right. \\
 &\quad \left. + 2t \frac{\sum_{i=1}^m |\alpha_i|}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_0^\infty |Nu(\tau)| d\tau \right) dr ds dt \right] \\
 &\leq e^{-t} t \left[\frac{1}{\|w\|_1^{p-1}} \|\alpha_r\|_1^{\frac{1}{p-1}} \right. \\
 &\quad \left. + 2 \|\alpha_r\|_1^{\frac{1}{p-1}} \frac{\sum_{i=1}^m |\alpha_i|}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) dr ds dt \right] \rightarrow 0 \text{ as } t \rightarrow \infty;
 \end{aligned}$$

$$\begin{aligned}
 e^{-t} |(Tu)'(t)| &\leq e^{-t} \left[\int_0^t \varphi_q \left(\frac{1}{w(r)} \right) \varphi_q \left(\int_r^\infty |Nu(\tau)| d\tau \right) dr \right. \\
 &\quad \left. + 2 \frac{\sum_{i=1}^m |\alpha_i|}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |Nu(\tau)| d\tau \right) dr ds dt \right] \\
 &\leq e^{-t} \left[\frac{1}{\|w\|_1^{p-1}} \|\alpha_r\|_1^{\frac{1}{p-1}} \right. \\
 &\quad \left. + 2 \|\alpha_r\|_1^{\frac{1}{p-1}} \frac{\sum_{i=1}^m |\alpha_i|}{|2 - \sum_{i=1}^m \alpha_i \xi_i^2|} \int_0^{\xi_i} \int_0^t \int_0^s \varphi_q \left(\frac{1}{|w(r)|} \right) dr ds dt \right] \rightarrow 0 \text{ as } t \rightarrow \infty,
 \end{aligned}$$

and

$$\begin{aligned}
 e^{-t} |(Tu)''(t)| &\leq e^{-t} \left[\varphi_q \left(\frac{1}{|w(r)|} \right) \varphi_q \left(\int_r^\infty |Nu(\tau)| d\tau \right) \right. \\
 &\quad \left. \leq e^{-t} \|\varphi_q \left(\frac{1}{w} \right)\|_\infty \|\alpha_r\|_1^{\frac{1}{p-1}} \rightarrow 0 \text{ as } t \rightarrow \infty. \right.
 \end{aligned}$$

This proves that $T(H)$ is equiconvergent at infinity. Therefore from lemma 2.1 we conclude that $T(H)$ is compact. The continuity of T can be proven using the Lebesgue dominated convergence theorem. This proves our lemma. \square

3. Existence result

We are now ready to prove our main existence Theorem.

Theorem 3.1 Let $K : [0, \infty) \times \mathfrak{R}^3 \rightarrow \mathfrak{R}$ be an $L^1[0, \infty)$ Caratheodory function. Suppose that there exist positive functions $a_i, b : [0, \infty) \rightarrow [0, \infty), a_i, b \in L^1[0, \infty), i = 1, 2, 3$ such that

$$|K(t, x, y, z)| \leq e^{-t(p-1)} [a_1(t) |u(t)|^{p-1} + a_2(t) |u'(t)|^{p-1} + a_3(t) |u''(t)|^{p-1}] + b(t)$$

a.e $t \in [0, \infty)$ with $D^{p-1} \sum_{i=1}^3 \|a_i\|_1 < 1$, where the constant D is defined in (2.21). Then the boundary value problem (1.1) - (1.2) has at least one solution for every $b \in L^1[0, \infty)$

Proof Let u be any solution of the λ dependent differential equation

$$(w(t)\varphi_p(u''(t)))' = \lambda K(t, u(t), u'(t), u''(t)) \text{ a.e } t \in [0, \infty), \lambda \in [0, 1] \tag{3.1}$$

Subject to the boundary condition (1.2)

Then

$$\begin{aligned} \|(w(t)\varphi_p(u''(t)))'\|_1 &= \lambda \|Nu\|_1 \\ &\leq \|a_1\|_1 \|u\|^{p-1} + \|a_2\|_1 \|u'\|^{p-1} + \|a_3\|_1 \|u''\|^{p-1} + \|b\|_1 \end{aligned}$$

From (2.17) we obtain

$$\begin{aligned} &\leq D^{p-1} \sum_{i=1}^3 \|a_i\|_1 \|h\|_1 + \|b\|_1 \\ &= D^{p-1} \sum_{i=1}^3 \|a_i\|_1 \|(w(t)\varphi_p(u''(t)))'\|_1 + \|b\|_1 \end{aligned}$$

which yields

$$\left(1 - \sum_{i=1}^3 \|a_i\|_1 D^{p-1}\right) \|(w(t)\varphi_p(u''(t)))'\|_1 \leq \|b\|_1$$

or

$$\|(w(t)\varphi_p(u''(t)))'\|_1 \leq \frac{\|b\|_1}{1 - D^{p-1} \sum_{i=1}^3 \|a_i\|_1}$$

and from (2.17) we obtain

$$\|u\| \leq D \|(w(t)\varphi_p(u''(t)))'\|_1^{\frac{1}{p-1}} \leq D \left(\frac{\|b\|_1}{1 - D^{p-1} \sum_{i=1}^3 \|a_i\|_1} \right)^{\frac{1}{p-1}}.$$

Therefore the set of solutions of (3.1)–(1.2) are a - priori bounded by a constant independent of λ and of solutions. The theorem is therefore proved. □

4. Example

Consider the boundary value problem

$$[w(t)\varphi_p(u''(t))] = e^{-2t} \left[\frac{|u(t)|^2}{(t+5)^2} + \frac{|u'(t)|^2}{10(t+1)^3} + \cos^2 t \frac{|u''(t)|^2}{10(t+1)^4} \right] + b(t) \tag{4.1}$$

$$u'(0) = \sum_{i=1}^2 \alpha_i \int_0^{\xi_i} u(t) dt, u(0) = 0, \lim_{t \rightarrow \infty} (w(t) \varphi_p(u''(t))) = 0 \tag{4.2}$$

$$w(t) = e^t, p = 3, q = \frac{3}{2}$$

where $b \in Z, b(t) > 0$

$$|K(t, u, u', u'')| = e^{-2t} \left[\frac{|u|}{(t+5)}, \frac{|u'|^2}{10(t+1)^3} + \cos^2 t \frac{|u''|^2}{10(t+1)^4} \right] + b(t)$$

$$a_1(t) = \frac{1}{(t+5)^2}, a_2(t) = \frac{1}{10(t+1)^3}, a_3(t) = \frac{1}{10(t+1)^4}$$

$$K(t, u, u', u'') \leq e^{-2t} \left[\frac{|u(t)|}{(t+5)} + \frac{|u'|^2}{10(t+1)^3} + \frac{|u''|^2}{10(t+1)^4} \right] + b(t)$$

$$\sum_{i=1}^3 \|a_i\|_1 = \frac{1}{5} + \frac{1}{20} + \frac{1}{30} = \frac{17}{60}$$

$$\alpha_1 = 8, \alpha_2 = 7, \xi_1 = \frac{1}{\sqrt{32}}, \xi_2 = \frac{1}{\sqrt{84}}$$

$$\sum_{i=1}^3 \alpha_i \xi_i^2 = \frac{1}{4} + \frac{1}{12} = \frac{1}{3} \neq 2$$

$$A = \frac{6}{5}e^{-1}, B = \frac{6}{5}, C = 1$$

$$D = \max(A, B, C) = \frac{6}{5}$$

$$D^{p-1} \sum_{i=1}^3 \|a_i\|_1 = \left(\frac{6}{5}\right)^2 \times \frac{17}{60} = \frac{51}{125} < 1$$

Therefore by Theorem 3.1 problem (4.1)-(4.2) has at least one solution for every $b \in Z$.

Acknowledgment

The authors would like to thank the Covenant University for providing financial support through grants during the study period.

Conflict of interest

The authors proclaim that there are no contending interests regarding the publication of this research paper.

Contribution of authors

The first author conceived the idea and developed the theory. The second author did the literature review. The results were discussed by all authors who also contributed to the final manuscript.

References

- [1] Agarwal RP, O' Regan D. Infinite interval problems for differential, difference and integral equations. Kluwer Academic, 2001.
- [2] Agarawal RP, O' Regan D. (Eds) Series in Mathematical analysis and applications. Vol. 3. *Gordon and Breach Science Publishers* (2001).
- [3] Frioui A, Guezane-Lakoud A, Khaldi A. Higher order boundary value problems at resonance on an unbounded interval. *Electronic Journal of Differential Equations* 2016; 4:1-10.
- [4] Gupta CP. A non-resonant multipoint boundary value problem of Neumann-Dirichlet type for a p -Laplacian type operator. *Dynamical Systems and Applications* 2004; 4: 439-442.
- [5] Hopkins B, Kosmatov N. Third-order boundary value problems with sign-changing solutions. *Nonlinear Analysis TMA* 2007; 67: 126-137.
- [6] Iyase SA. On a third-order boundary value problem at resonance on the half-line. *Araian Journal of Mathematics* 2019; 8: 43-53. doi: 202007 540065 – 018-0209-5
- [7] Kim CG. Solvability of multipoint boundary value problems on the half-line. *Journal of Nonlinear Science and Applications* 2012; 5: 27-33.
- [8] Kosmatov N. Second order boundary value problems on an unbounded domain. *Nonlinear Analysis TMA* 2008; 68: 2158-2171.
- [9] Lian H, Ge W. Solvability of second order three point boundary value problems on the half-line. *Applied Mathematical Letters* 2003; 16: 33-39.
- [10] O'Regan D, Precup R. Theorems of Leray-Schauder type and applications. CRC Press, 2002.
- [11] Zeidler E. *Nonlinear functional analysis and its applications*. Springer-Verlag, New York, 1986.