

- [JOURNALS](#)
- [CONFERENCES](#)
- [EBOOKS](#)
- [PUBLISHERS](#)
- [ABOUT](#)
- [CITATION MANAGER](#)

- [HOME](#)
- [LATEST CONFERENCE](#)
- [ALL YEARS](#)
- [OTHER PROCEEDINGS](#)
- [VISIT SPE](#)
- [CITATION MANAGER](#)

## **SPE Nigeria Annual International Conference and Exhibition**

August 11–13, 2020

Virtual

[\*\*Day 1 Tue, August 11, 2020\*\*](#)

ISBN:

978-1-61399-785-7

- [Previous Paper](#)
- [Next Paper](#)

# **Experimental Studies on the Performance of Bio Based and Industrial Surfactants in Enhanced Oil Recovery**

[Damilola Abraham;](#)

[Oyinkepreye Orodu;](#)

[Vincent Efeovbokhan;](#)

[Emmanuel Okoro;](#)

[Temiloluwa Ojo;](#)

[Lekan Keshinro](#)

Paper presented at the SPE Nigeria Annual International Conference and Exhibition, Virtual, August 2020.

Paper Number: SPE-203759-MS

<https://doi.org/10.2118/203759-MS>

Published: August 11 2020

- **Cite**

- [Share Icon](#)**Share**
- **Get Permissions**

## Abstract

Surfactants are known for their unique property in lowering the interfacial tension (IFT) amid fluids injected and heavy crude oil of 22.3°API. In this present work, an original surfactant was formulated from natural oil (Castor oil) to see its use in enhanced oil recovery. The results from interfacial tension reduction by the castor oil-based surfactant were compared to that of the industrial surfactant (Methyl ester sulfonate). The IFT between the aqueous phases was measured then the effect of the surfactants was studied in core flooding experiments. The IFT was found to be reduced to as low as 12.1 mN/m using the castor-based surfactant and 12.3 mN/m using the industrial surfactant. The influence of brine concentration on IFT was also investigated. Results from core floods showed that the range of oil recovery after waterflood is in the range of 30-40% and the additional recovery from surfactant flooding in the range of 35-46%.

### Keywords:

[enhanced recovery](#), [ift](#), [brine concentration](#), [crude oil](#), [waterflooding](#), [concentration](#), [surfactant flooding](#), [chemical flooding methods](#), [industrial surfactant](#), [methyl ester sulfonate](#)

### Subjects:

[Improved and Enhanced Recovery](#), [Waterflooding](#), [Chemical flooding methods](#)

Copyright 2020, Society of Petroleum Engineers

You can access this article if you purchase or spend a download.

## Sign in

[Reset password](#)

[Register](#)

[Sign in via OpenAthens](#)

Pay-Per-View Access

\$28.00

[BUY THIS ARTICLE](#)

**Annual Article Package – 25**

\$225

[BUY DOWNLOADS](#)

**Annual Article Package – 50**

\$400

[BUY DOWNLOADS](#)

[View Your Downloads](#)

**SPE members can access this article  
for \$7 USD. [Learn how to connect your  
SPE membership.](#)**

[View Metrics](#)

### **Email Alerts**

[Proceedings Paper Activity Alert](#) [Alert](#)

[Latest Conference Proceeding Alert](#)

Advertisement

### **Suggested Reading**

[Effect of Time and Temperature on Crude Oil Aging to do a Right Surfactant Flooding with a New Approach](#)

14OTCA

[Enhanced Oil Recovery for Carbonate Oil Reservoir by using Nano-Surfactant:Part I](#)

19GOTS

[Application of Sodium Lauryl Sulfate for Enhanced Oil Recovery of Medium Crude Oil in the Niger Delta Fields](#)

22NAIC

[Non-Edible Oil Based Surfactant For Enhanced Oil Recovery](#)

21NAIC

[NanoSurfactant for EOR in Carbonate Reservoirs](#)

17SATS