




BJPT British Journal of Pharmacology and Toxicology



ISSN (Online): 2044-2467
ISSN (Print): 2044-2459

[Submit an article](#)

Current Information

- Paper in Press
- Current Issue
- Archives
- Author's Comments

Sales & Services

- Subscribe / Renew
- Recommend to Library
- Reprints/Offprints
- Special Sale
- Customer Services

Contact Information

Executive Managing Editor
Email: admin@maxwellsci.com
Publishing Editor
Email: support@maxwellsci.com
Account Manager
Email: faisalm@maxwellsci.com
Journal Editor
Email: admin@maxwellsci.com
Press Department
Email: press@maxwellsci.com

Abstract

2011 (Vol. 2, Issue: 3)

Article Information:

Toxicity Evaluation of the Liver and *in vitro* Metabolism in Wistar Rat on Exposure to N-Nitrosamine Precursors

Usunobun Usunomena, Josiah J. Sunday, Nwangwu Spencer, Uhunmwagho S. Esosa, Omage Kingsley and Maduagwu N. Emmanuel

Corresponding Author: Usunobun Usunomena

Key words: Dimethylamine hydrochloride, incubation, N-nitrosamine, Sodium nitrite, UV irradiation, , Vol. 2 , (3): 138-142

Submitted	Accepted	Published
2011 May, 23	2011 July, 02	2011 August, 05

Abstract:

The aim of this study is to evaluate liver toxicity on exposure to n-nitrosamine precursors as well as the effect of ultraviolet light on n-nitrosamines and its precursors. Toxicological evaluation of the liver following single dose treatment of wistar rat with 8.2125 mg NaNO₂/adult rat and in rats given a combined dose of 50 mgDMA-HCl and 8.2125 mg NaNO₂/adult rat showed a steady elevation of the liver function enzymes. Histopathological analysis of the liver showed hepatic necrosis in the chemical induced rats. Following UV exposure after *in vitro* incubation of rat liver microsomal plus soluble fraction with NaNO₂ and NaNO₂ plus DMA-HCl, nitrite concentration in the NaNO₂ incubation medium was 19.5 and 2.2 µg/mL before and after UV exposure respectively while the nitrite concentration in the NaNO₂ plus DMA-HCl incubation medium was 23.5 and 2.5 µg/mL, respectively. Nitrite loss was significant (p<0.05) between before and after UV exposure in all groups. UV exposure, thus degraded the nitrosamine precursors, nitrite and DMA-HCl, thereby inhibiting possible nitrosation. The high values of the activities of serum transaminases (AST and ALT), alkaline phosphatases (ALP) and gamma-glutamyltransferses (γ-GT), relative to control values are indicative of severe intrahepatic cell damage.

Abstract	PDF	HTML
Abstract	PDF	HTML

Cite this Reference:

Usunobun Usunomena, Josiah J. Sunday, Nwangwu Spencer, Uhunmwagho S. Esosa, Omage Kingsley and Maduagwu N. Emmanuel, 2011. Toxicity Evaluation of the Liver and *in vitro* Metabolism in Wistar Rat on Exposure to N-Nitrosamine Precursors. British Journal of Pharmacology and Toxicology, 2(3): 138-142.

Advertise with us