




Available online at <http://www.urpjournals.com>

## International Journal of Agricultural and Food Science

Universal Research Publications. All rights reserved



### Original Article

ISSN 2249-8516

#### Microbiological safety evaluation of snacks sold in fast food shops in Ota, Ogun state, Nigeria.

\*Oranusi, S; Omagbemi, F. and Eni, A.O.

Department of Biological Sciences, Covenant University, Canaanland, Ota, Ogun State, Nigeria.

\*Correspondent: orasol2002@yahoo.com.au

Received 28 October 2011; accepted 08 November 2011

#### Abstract

The microbial quality of snacks (ready to eat foods) sold in Ota, Ogun State was investigated. A total of 100 different samples from 3 vending sites namely, a University Cafeteria, a top class snacks bar and a local kiosk were analyzed for total aerobic plate count, coliform count and for specific pathogens and fungi. The University Cafeteria had mean total aerobic plate count and coliform count ranging from  $1.1 \times 10^3$ - $3.0 \times 10^4$  and  $1.0 \times 10^2$ - $2.2 \times 10^3$ . The snacks bar had mean total aerobic plate count and coliform count ranging from  $2.0 \times 10^3$ - $5.8 \times 10^5$  and  $1.4 \times 10^2$  - $1.8 \times 10^5$  while the local kiosk had mean total aerobic plate count and coliform count ranging from  $2.1 \times 10^3$ - $5.4 \times 10^5$  and  $1.0 \times 10^2$  - $8.0 \times 10^4$  respectively. The fungal counts from the three sites are within  $1.0 \times 10^2$ -  $4.0 \times 10^2$ . Six different bacterial and three fungal isolates were identified to include *E. coli*, *S. aureus*, *Bacillus cereus*, *Enterococcus*, *Klebsiella spp*, *Pseudomonas spp* and *Aspergillus niger*, *Penicillium spp* and *Mucor*. The presence of *E. coli* and Enterococci which are indicator organisms call for concern. Adoption of good manufacturing practice and hazard analysis critical control point (HACCP) are necessary to preventing occurrence of food borne illness.

© 2011 Universal Research Publications. All rights reserved

**Key words:** Food safety, Pathogens, Snacks, Ready to eat foods, Coliform.

#### 1. Introduction

Food borne illnesses are diseases, usually either infectious or toxic in nature, caused by agents that enter the body through the ingestion of food [1]. Microbial agents that cause food borne illness may include, bacteria such as; *Salmonella*, *Staphylococcus aureus*, *Escherichia coli* (pathogenic strains) *Bacillus spp*, *Clostridium botulinum*, *Listeria monocytogens*; viruses such as; hepatitis A and E, Norovirus; molds, fungi and yeasts [2]. In addition, poisonous chemicals, or other harmful substances can cause food borne diseases if they are present in food. Symptoms of food borne illnesses may differ amongst pathogens but general symptoms may include diarrhea, nausea, vomiting, fever, and abdominal cramps. Some can cause organ failure.

Ready to eat foods are foods that are consumed in the same state as that in which it is sold and does not include nuts in the shell and whole, raw fruits and vegetables that are intended for hulling, peeling or washing by the consumer [3]. Some ready-to-eat foods are regarded as potentially hazardous, because such foods can support the growth of pathogens. Such food must be kept at certain temperatures and conditions to minimise the growth of any pathogens that

may be present in the food or to prevent the formation of toxins in the food. There is a wide variety of ready-to-eat foods, including, but are not limited to sandwiches, kebabs, hotdogs, meat pie, salad, doughnuts, takeaway foods and bakery products. Ready-to-eat foods usually include a number of ingredients which may or may not be cooked. Due to the nature of these foods and their methods of preparation involving extensive handling, they are usually prone to contamination/cross contamination from soil, water, air, storage/distribution facilities, environment and human activities (food handlers and vendors).

Convenience /modern life style, industrialization, economic down turn, quest for more wealth, materialism, and their associated lack of time to prepare proper meal, low purchasing power, are reasons for the increased patronage for ready to eat foods [4].

WHO [1] estimated that a significant proportion of the approximately 1.5 billion episodes of diarrhea and more than 3 million deaths globally recorded annually, results from consumption of food with microbial pathogens and toxins. We live in a microbial world, and there are many

opportunities for food to become contaminated as it is produced and prepared, many of these food borne microbes are found in healthy animals raised for food (usually in their intestines). Meat and poultry carcasses can become contaminated during slaughter by contact with small amounts of intestinal contents. Similarly, fresh fruits and vegetables can be contaminated if they are washed or irrigated with water that is contaminated with animal manure or human sewage. A number of foods in Nigeria have been reported to have high level of contaminants [5, 6, 7, 8, 9, 10], however, there is scanty information about the extent of microbial contamination of snacks sold in Nigerian fast food shops and supermarkets. This study was undertaken to determine the microbial profile of snacks sold amongst 3 socio economic classes (upper, middle and lower class) in Ota, Ogun state, Nigeria with a view of proffering food safety advice.

## 2. Materials and Methods

### 2.1 Sources of sample

Three ready-to eat-food vending sites in Ota, Ogun state were sampled. These sites were chosen because they are highly patronized by members of the general public of different socio economic classes (upper, middle and lower class). These locations include; a University cafeteria (upper class), a snacks bar/fast food shop (middle class) and a local kiosk (lower class).

### 2.2 Sample collection

Twenty (20) samples each of fresh Hot dog, Sausage, Meat pie, Egg roll and Doughnuts were purchased from these locations. Samples were randomly selected without order. Two each of the samples were collected on alternate days from the sites. The samples were aseptically collected in sterile polyethylene bags and transferred immediately to the laboratory for further analysis.

### 2.3 Culturing of food samples

Ten (10) gram of each food samples was blended and homogenized in 100 ml of sterile distilled water ( $10^{-1}$  dilution). Serial dilutions of the homogenates were made to  $10^{-2}$  and  $10^{-3}$ . One (1) ml of each dilution was plated in replicate using both pour plate and spread plate methods on Nutrient agar for total aerobic plate count and isolation of other microorganisms, Eosin methylene blue agar for coliform count and isolation. Mannitol salt agar and Potato Dextrose agar (PDA) plus gentamycin were inoculated for Staphylococci and fungi isolation. The plates were incubated at  $37^{\circ}\text{C}$  for 24hrs except for PDA that was incubated for 3-5 days at room temperature  $28-30^{\circ}\text{C}$ .

### 2.4 Coliform test

**Presumptive test:** One (1) gram of each sample was transferred to sterile McCartney bottles containing Lactose broth and inverted Durham tubes. Incubation was for 24-48hrs at  $37^{\circ}\text{C}$ . Tubes showing gas production and/or colour change of dye were streaked on EMB plates. Incubation of plates for **Confirmatory test** was at  $37^{\circ}\text{C}$  and  $44^{\circ}\text{C}$  for 24hrs;

colonies from EMB plates were picked and inoculated into tubes containing lactose broth for **completed test** and onto Nutrient agar slants for further characterization. Inoculated tubes and slants were incubated for 24hrs at  $37^{\circ}\text{C}$  [11].

### 2.5 Identification of isolates

The bacteria isolates were identified based on standard methods [12, 13]. Fungal isolates were identified based on cultural and morphological characteristics with reference to standard atlas [14].

### 2.6 Statistical analysis

The values obtained for total aerobic and coliform counts were subjected to analysis of variance [15].

## 3. Results

The mean total aerobic plate count of sample from the vending sites is as shown in Table 1 It shows that hot dog from local kiosk and egg roll from snacks bar had significant higher counts compared to other samples. It also reveals that samples from the local kiosk and snacks bar had relatively higher counts compared to the University cafeteria.




Fig.1. Percentage occurrence of bacterial isolates from samples




Fig.2. Percentage occurrence of fungal isolates from samples

**Table 1.** Mean Total Aerobic Plate Count (cfu/g)

| Food sample | Source of samples                  |                                    |                                     |
|-------------|------------------------------------|------------------------------------|-------------------------------------|
|             | University cafeteria               | Snacks bar                         | Local kiosk                         |
| MEAT PIE    | 2.0x 10 <sup>3</sup> <sub>a</sub>  | 2.7x10 <sup>4</sup> <sub>a</sub>   | 2.1 x 10 <sup>3</sup> <sub>a</sub>  |
| HOTDOG      | 2.0x 10 <sup>3</sup> <sub>a</sub>  | 2.8x10 <sup>3</sup> <sub>a</sub>   | 5.4 x 10 <sup>5</sup> <sub>b</sub>  |
| SAUSAGE     | 3.0x10 <sup>4</sup> <sub>a</sub>   | 3.3 x 10 <sup>4</sup> <sub>a</sub> | 8.0 x 10 <sup>3</sup> <sub>a</sub>  |
| EGG ROLL    | 8.2x10 <sup>3</sup> <sub>a</sub>   | 5.8 x 10 <sup>5</sup> <sub>b</sub> | 4.8 x 10 <sup>4</sup> <sub>ab</sub> |
| DOUGHNUT    | 1.1 x 10 <sup>3</sup> <sub>a</sub> | 2.0 x 10 <sup>3</sup> <sub>a</sub> | 3.0 x 10 <sup>4</sup> <sub>a</sub>  |

a, b= Mean within row with the same letter for same count are not significantly different (p>0.05)

**Table 2.** Mean Coliform Count (cfu/g)

| Food sample | Source of samples                  |                                    |                                    |
|-------------|------------------------------------|------------------------------------|------------------------------------|
|             | University cafeteria               | Snacks bar                         | Local kiosk                        |
| MEAT PIE    | NG                                 | 6.0 x 10 <sup>2</sup> <sub>a</sub> | 2.0 x 10 <sup>2</sup> <sub>a</sub> |
| HOTDOG      | 1.0 x 10 <sup>2</sup> <sub>a</sub> | 1.4 x 10 <sup>2</sup> <sub>a</sub> | 5.0 x 10 <sup>2</sup> <sub>a</sub> |
| SAUSAGE     | 2.2 x 10 <sup>3</sup> <sub>a</sub> | 1.8 x 10 <sup>5</sup> <sub>b</sub> | 1.0 x 10 <sup>3</sup> <sub>a</sub> |
| EGG ROLL    | 6.0 x 10 <sup>2</sup> <sub>a</sub> | 1.0 x 10 <sup>3</sup> <sub>a</sub> | 8.0 x 10 <sup>4</sup> <sub>a</sub> |
| DOUGHNUT    | NG                                 | NG                                 | 1.0 x 10 <sup>2</sup> <sub>a</sub> |

a, b= Mean within row with the same letter for same count are not significantly different (p>0.05)

**Table 3.** Mean Fungal Count (cfu/g)

| Food sample | Source of samples    |                      |                     |
|-------------|----------------------|----------------------|---------------------|
|             | University cafeteria | Snacks bar           | Local kiosk         |
| MEAT PIE    | NG                   | 3.0 x10 <sup>2</sup> | NG                  |
| HOTDOG      | NG                   | 2.0x10 <sup>2</sup>  | 2.0x10 <sup>2</sup> |
| SAUSAGE     | 4.0x10 <sup>2</sup>  | NG                   | 3.0x10 <sup>2</sup> |
| EGG ROLL    | 3.4x10 <sup>2</sup>  | 2.8x10 <sup>2</sup>  | 4.0x10 <sup>2</sup> |
| DOUGHNUT    | 2.0x10 <sup>2</sup>  | 1.0x10 <sup>2</sup>  | NG                  |

Table 2 Shows mean coliform count of samples; it reveals that sausage from snacks bar had significant higher counts than sausage from other sites. Table 3 Shows mean fungal count of samples from the three sites. Egg roll had growth in all the cases. *E. coli*, *S. aureus*, *Pseudomonas spp*, *Enterococcus*, *Bacillus cereus*, *klebsiella spp*, *Aspergillus niger*, *Mucor* and *Penicillium* were isolated as shown in Figs.1 and 2

#### 4. Discussion

The snacks sold in these vending sites had microbial loads within acceptable microbiological quality, ICMSF [16]; Food

Standards [17], declares ready to eat foods with aerobic plate counts 10<sup>4</sup> - < 10<sup>5</sup> as acceptable. Egg roll and Sausage had higher plate counts, these could be due to the nutritional content of the major ingredient used in their preparation i.e. eggs and hams, which offers a rich nutrient media for microbial growth [18]. The presence of coliforms points to poor sanitary practices by food personnel and could be an indication of possible fecal contamination. Doughnut preparation does not involve egg and meat addition, this could have accounted for the relatively lower count for this product. The higher counts in microbial loads observed in the

local kiosk and snack bar could be attributed to the levels of exposure of these products.

The presence of *E. coli*, *Enterococcus*, *Klebsiella spp*, *Staphylococcus aureus*, is of concern and further support the possibility of fecal contamination of products due to poor sanitation [19].

*Bacillus* and *Pseudomonas spp* were isolated; however, *Salmonella* and *Shigella spp* were not detected. These organisms are known to be environmental contaminants and opportunistic pathogens [10], have been implicated in food borne diseases, and are known to cause food spoilage that can lead to economic loss. The most predominant bacterial contaminants was *S. aureus* with 25.56% this could be traced to the fact that it is abundant in human body (skin, nails hair) [20, 21, 22]. Similarly, *Bacillus cereus* showed high percentage (18%), its presence can be traced to the fact that it is abundant spore former in soil, air and water, hence can easily be present in these foods. This report is in agreement to reports of [6, 8, 9], they isolated similar organisms from sausages, meat pie and sea foods respectively.

The presence of *Aspergillus*, *Penicillium* and *Mucor* could be attributed to the surrounding air and packaging materials [23, 24]. *Aspergillus spp* are very common fungal agent of food borne illness, [25, 26]. A comparison of the level of contamination of snacks from sampling points representing 3 socio-economic classes: upper class, middle class and lower class shows no significant difference in their levels of contamination, although the snacks bar and the local kiosk had relatively higher level of contamination compared to the university cafeteria

Snacks (ready to eat foods) are eaten by all age groups with high popularity amongst school children and youths, it is therefore mandatory that these foods must be free from contamination as much as possible. Food borne illness can be prevented by good hygiene practice during the preparation of food. To prevent occurrence of food borne illness it is therefore, important to ensure that foods sold are safe and hygienic, public awareness programs should be employed to educate personnel involved in food preparation, food processors, and food vendors. The general public should be educated on the need for food safety and the requirement for water meant for human consumption and for food processing [27, 28, 29]. Proper and regular hand washing, sanitization of all equipment and utensils, care for the environment and the packaging materials so as to prevent the spread of contaminants will help in safety of food. Adoption of the HACCP (Hazard Analysis Critical Control Point) principle in snacks preparation is advocated.

## References

- [1] WHO, Food Safety and Foodborne Illness. Fact Sheet 237 Review: World Health Organization, Geneva, Switzerland (2007).
- [2] CDC, Preliminary Food Net data on the incidence of infection with pathogens transmitted commonly through food. *Morb Mortal Wkly Rep.* 59(14) (2010) 418-22
- [3] R.J. Gilbert, J. De-Louvois, T. Donovan, C. Little, K. Nye, Guidelines for the microbiological quality of some ready-to-eat foods samples at the point of ne sale. *Communicable Disease and Public Health* 3(2000) 163-167.
- [4] A.C. Nielsen, Ready to eat meals: Global highlights. A global ACNielsen consumer survey report (2006)
- [5] A.A. Adesiyun, Bacteriologic quality of some Trinidadian ready to consume foods and drinks and possible health risks to consumers. *J. Food Prot.* 58(3) (1995) 651-655.
- [6] F. Oluwafemi, M.T. Simisaye, Extent of Microbial Contamination of Sausages sold in two Nigerian cities. *Afri. J. Biomed. Res.* 9 (2005) 133 – 136
- [7] U. Shamsuddeen, Microbiological quality of spice used in the production of Kilishi a traditionally dried and grilled meat product. *Bajopas.* 2(2) (2009) 66-69
- [8] S.C. Yah, C.O. Nwinyi, N.S. Chinedu, Assessment of bacteriological quality of ready to eat food (Meat pie) in Benin City metropolis, Nigeria. *African Journal of Microbiology Research.* 3(6) (2009) 390-395
- [9] I.O. Okonko, O.D. Adejoye, A.A. Ogun, A.A. Ogunjobi, A.O. Nkang, B.C. Adebayo, Hazards analysis critical control points (HACCP) and microbiology qualities of sea-foods as affected by handler's hygiene in Ibadan and Lagos, Nigeria. *Afr. J. Food Sci.* 3(2) (2009) 35-50
- [10] A.O. Eni, A.O. Ibukunoluwa, U.S. Oranusi, Microbial quality of fruits and vegetables sold in Sango Ota, Nigeria. *Afr. J. Food Sci.* 4(5) (2010) 291-296.
- [11] S. Oranusi, E. Onyeike, M. Galadima, V.J. Umoh, Hazard analysis critical control points of foods prepared by families in Zaria, Nigeria. *Nig. J. Microbiol.* 18(1-2) (2004) 346-362
- [12] M.L. Speck, Compendium of Methods for Microbiological Examination of Foods. American Public Health Association, Washington D.C. (1976) Pp 277-328
- [13] M. Cheesbrough, District Laboratory Practice in Tropical Countries Part I. Cambridge University press, United Kingdom (2004) Pp 178-209
- [14] W. Tsuneo, Pictorial atlas of soil and seed fungi: Morphologies of cultural fungi and key to species. Third editon. CRC press (2010).

- [15] G.W. Snedecor, W.C. Cochran, Statistical methods. The Iowa State College Press, Ames, Iowa, USA (1976).
- [16] ICMSF (International Commission on Microbiological Specification for Food), Sampling for microbiological analysis. Principles and specific application. University of Toronto press, Toronto (1974). Pp 1-18
- [17] Food Standards, Australia-New Zealand Guidelines for the microbiological examination of ready-to-eat foods. Retrieved Sept. 10 2010 from, [http://www.foodstandards.gov.au/\\_srcfiles/Guidelines%20for%20Micro%20exam.pdf](http://www.foodstandards.gov.au/_srcfiles/Guidelines%20for%20Micro%20exam.pdf).
- [18] M. Phillips, Analysis of Microbial Hazards related to time/temperature control of foods for safety. *Comprehensive Review in Food Science and Food safety*. 2(2003) 33-35
- [19] S. Oranusi, M. Galadima, V.J. Umoh, P.I. Nwanze, Food safety evaluation in boarding schools in Zaria, Nigeria, using the HACCP system. *Sc. Res. Essays*. 2(10) (2007) 426-433
- [20] N. Balaban, A. Rasooly, Staphylococcal enterotoxins. *Int. J. Food Microbiol.* 61(2000) 1-10
- [21] S. Oranusi, M. Galadima, V.J. Umoh, Toxicity test and bacteriophage typing of *S. aureus* isolates from food contact surfaces and foods prepared by families in Zaria, Nigeria. *Afr. J. Biotechnol.* 5(4) (2006a) 362-365
- [22] S. Oranusi, M. Galadima, V.J. Umoh, Phage typing and toxigenicity test of *S. aureus* strains from food contact surfaces and foods prepared in boarding schools in Zaria, Nigeria. *Nig. J. Microbiol.* 20(2) (2006b) 1011-1017
- [23] R.I. Aboloma, Microbiological analysis of bread samples from bakery to sale points in Ado-Ekiti, Ekiti State. Nigeria. *Biological and Environmental Sciences Journal for the Tropics* 5(3) (2008) 77-81.
- [24] A.H. Kawo, F.N. Abdulmumin, Microbiological quality of re-packaged Sweets sold in metropolitan Kano, Nigeria. *Bajopas*. 2(1) (2009) 154 – 159
- [25] M. Peraica, A.M. Domijan, Mycotoxins in food and human health. *Arh Hig. Rada Toksikol* 52 (2001) 23-35
- [26] S. Katherine, M. Catherine, F. Rachel, Mycotoxins explained. *Food Safety & Hygiene*. A bulletin for the Australian food industry (2006).
- [27] S. Taulo, A. Wetlesen, R. Abrahamsen, R. Mkakosya, G. Kululanga, Microbiological quality of water, associated management practices and risks at source, transport and storage points in a rural community of Lungwena, Malawi. *Afr. J. Microbiol. Res.* 7(2) (2008) 131-137
- [28] I. O. Okonko, O.D. Adejoye, T.A. Ogunnusi, E.A. Fajobi, O.B. Shittu, Microbiological and physicochemical analysis of different water samples used for domestic purposes in Abeokuta and Ojota. Lagos State, Nigeria. *Afr. J. Biotechnol.* 7(3) (2008a) 617-621
- [29] I.O. Okonko, A.A. Ogunjobi, O.D. Adejoye, T.A. Ogunnusi, M.C. Olasogba, Comparative studies and Microbial risk assessment of different water samples used for processing frozen sea-foods in Ijoraolopa, Lagos State, Nigeria. *Afr. J. Biotechnol.* 7(16) (2008b) 2902 2907

Source of support: Nil; Conflict of interest: None declared