



ELSEVIER

## Biochemical Pharmacology

Volume 32, Issue 23, 1 December 1983, Pages 3577-3581



# Nitrosamine metabolism in kwashiorkor rats

[Emmanuel N. Maduagwu. Author links open the author workspace.\\*. Numbers and letters correspond to the affiliation list. Click to expose these in author workspace](#) [Eva Frei. Author links open the author workspace.](#) [Norbert Frank. Author links open the author workspace.](#) [Bertold Spiegelhalter. Author links open the author workspace.](#) [Rolf Preussmann. Author links open the author workspace.](#)

Institute of Toxicology and Chemotherapy, German Cancer Research Center, Im Neuenheimer Feld 280, D-6900 Heidelberg, F.R.G.

[http://dx.doi.org/10.1016/0006-2952\(83\)90306-4](http://dx.doi.org/10.1016/0006-2952(83)90306-4) [Get rights and content](#)

## Abstract

The *in vitro* metabolism of nitrosodimethylamine (NDMA) was studied in liver tissue obtained from male weanling kwashiorkor wistar rats. The elimination of this compound and that of nitrosomorpholine (NMOR) from the blood, after a single intravenous dose, was also investigated. *N*-demethylase activity in liver microsomes of the test animals was not significantly different from that of the controls although the activity of this enzyme per gram wet liver tissue was considerably reduced in the model animals. On the other hand, the glutathione (GSH) content in liver cytosol of the kwashiorkor animals was much higher than that of the controls. The elimination of NDMA and NMOR from the blood of the experimental animals over 8 hr following i.v. administration of the carcinogens, showed that the clearance rate of each nitrosamine was significantly lower in the kwashiorkor rats.

Copyright © 1983 Published by Elsevier Inc.