

Research Article

An Extension of Gregus Fixed Point Theorem

J. O. Olaleru and H. Akewe

Received 2 October 2006; Accepted 17 December 2006

Recommended by Lech Gorniewicz

Let C be a closed convex subset of a complete metrizable topological vector space (X, d) and $T : C \rightarrow C$ a mapping that satisfies $d(Tx, Ty) \leq ad(x, y) + bd(x, Tx) + cd(y, Ty) + ed(y, Tx) + fd(x, Ty)$ for all $x, y \in C$, where $0 < a < 1$, $b \geq 0$, $c \geq 0$, $e \geq 0$, $f \geq 0$, and $a + b + c + e + f = 1$. Then T has a unique fixed point. The above theorem, which is a generalization and an extension of the results of several authors, is proved in this paper. In addition, we use the Mann iteration to approximate the fixed point of T .

Copyright © 2007 J. O. Olaleru and H. Akewe. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Gregus [1] proved the following theorem.

THEOREM 1.1. *Let C be a closed convex subset of a Banach space X and $T : C \rightarrow C$ a mapping that satisfies $\|Tx - Ty\| \leq a\|x - y\| + b\|x - Tx\| + c\|y - Ty\|$ for all $x, y \in C$, where $0 < a < 1$, $b \geq 0$, $c \geq 0$, and $a + b + c = 1$. Then T has a unique fixed point.*

Several papers have been written on the Gregus fixed point theorem. For example, see [2, 3]. The theorem has been generalized to the condition when X is a complete metrizable topological vector space [4].

When $a = 1$, $b = 0$, $c = 0$, T becomes a nonexpansive map. In the past four decades, several papers have been written on the existence of a fixed point (which may not be unique) for a nonexpansive map defined on a closed bounded and convex subset C of a Banach space X . For example, see [5–7]. Recently, the existence of fixed points of T when the domain of T is unbounded was discussed in [6]. When $a = 0$, we have the Kannan maps. Similarly, several papers have been written on the existence of a fixed point for a

2 Fixed Point Theory and Applications

Kannan map defined on a Banach space, for example, see [8, 9]. The fixed point theorem of Gregus is interesting because it tells what happens if $0 < a < 1$.

Chatterjea [10] considered the existence of fixed point for T when T is defined on a metric space (X, d) , such that for $0 < a < 1/2$,

$$d(Tx, Ty) \leq a\{d(x, f(y)) + d(y, f(x))\}. \quad (1.1)$$

It is natural to combine this condition with that of Gregus to get the following condition:

$$d(Tx, Ty) \leq ad(x, y) + bd(x, Tx) + cd(y, Ty) + ed(y, Tx) + fd(x, Ty) \quad (1.2)$$

for all $x, y \in C$, where $0 < a < 1$, $b \geq 0$, $c \geq 0$, $e \geq 0$, $f \geq 0$, and $a + b + c + e + f = 1$.

Observe that if T satisfies (1.2), then it also satisfies

$$d(Tx, Ty) \leq ad(x, y) + pd(x, Tx) + pd(y, Ty) + pd(y, Tx) + pd(x, Ty) \quad (1.3)$$

for all $x, y \in C$, where $0 < a < 1$, $p \geq 0$, $a + 4p = 1$, ($p = (1/4)b + (1/4)c + (1/4)e + (1/4)f$). Thus b , c , e , and f will be used interchangeably as p in the proof of our main theorem.

As observed by Chidume [5, page 119], since the four points $\{x, y, Tx, Ty\}$ of (1.2) determine six distances in X , the inequality amounts to say that the image distance $d(Tx, Ty)$ never exceeds a fixed convex combination of the remaining five distances. Geometrically, this type of condition is quite natural.

In this paper, we extend Gregus result to the condition when T satisfies condition (1.2) and also generalize it to the condition when X is a complete metrizable topological vector space, thus answering the question posed in [4]. Complete metrizable topological vector spaces include uniformly convex Banach spaces, Banach spaces and complete metrizable locally convex spaces (see [11, 12]).

The following result will be needed for our result.

THEOREM 1.2 [13, 14]. *A topological vector space X is metrizable if and only if it has a countable base of neighbourhoods of zero. The topology of a metrizable topological vector space can always be defined by a real-valued function $\|\cdot\| : X \rightarrow \mathfrak{R}$, called F -norm such that for all $x, y \in X$,*

- (1) $\|x\| \geq 0$,
- (2) $\|x\| = 0 \Rightarrow x = 0$,
- (3) $\|x + y\| \leq \|x\| + \|y\|$,
- (4) $\|\lambda x\| \leq \|x\|$ for all $\lambda \in K$ with $|\lambda| \leq 1$,
- (5) if $\lambda_n \rightarrow 0$, and $\lambda_n \in K$, then $\|\lambda_n x\| \rightarrow 0$.

For the same result see Kothe [15, Section 15.11]. Henceforth, unless otherwise indicated, F will denote an F -norm if it is characterizing a metrizable topological vector space. Observe that an F -norm will be a norm if it is defining a normed space.

We now prove our main theorem. We use the technique in [4] which is due to Gregus [1].

THEOREM 1.3. *Let C be a closed convex subset of a complete metrizable space X and $T : C \rightarrow C$ a mapping that satisfies $F(Tx - Ty) \leq aF(x - y) + bF(x - Tx) + cF(y - Ty) + eF(y - Tx) + fF(x - Ty)$ for all $x, y \in C$, where $0 < a < 1$, $b \geq 0$, $c \geq 0$, $e \geq 0$, $f \geq 0$, and $a + b + c + e + f = 1$. Then T has a unique fixed point.*

Proof. Take any point $x \in C$ and consider the sequence $\{T_n(x)\}_{n=1}^\infty$,

$$\begin{aligned} F(T^n x - T^{n-1} x) &\leq aF(T^{n-1} x - T^{n-2} x) + bF(T^{n-1} x - T^n x) \\ &\quad + cF(T^{n-2} x - T^{n-1} x) + eF(T^{n-2} x - T^n x) \\ &\quad + fF(T^{n-1} x - T^{n-1} x) \\ &\leq \frac{a+c+e}{1-b-e} F(T^{n-1} x - T^{n-2} x) \\ &\leq \frac{a+2p}{1-2p} F(T^{n-1} x - T^{n-2} x) \leq F(Tx - x). \end{aligned} \tag{1.4}$$

Thus

$$F(T^n x - T^{n-1} x) \leq F(Tx - x). \tag{1.5}$$

In effect, it means that the distance between two consecutive elements of $\{T^n x\}$ is less or equal to the distance between the first and the second element. Now let us consider the distance between two consecutive elements with odd (resp., even) power of T . It is sufficient to consider only the distance between Tx and T^3x ,

$$\begin{aligned} F(T^3 x - Tx) &\leq aF(T^2 x - x) + bF(T^2 x - T^3 x) + cF(Tx - x) \\ &\quad + eF(x - T^3 x) + fF(T^2 x - Tx) \\ &\leq aF(T^2 x - Tx) + aF(Tx - x) + bF(T^2 x - T^3 x) \\ &\quad + cF(Tx - x) + eF(x - Tx) + eF(Tx - T^2 x) \\ &\quad + eF(T^2 x - T^3 x) + fF(T^2 x - Tx) \\ &\leq (2a + b + c + 3e + f)F(Tx - x) = (a + 2p + 1)F(Tx - x). \end{aligned} \tag{1.6}$$

Hence

$$F(T^3 x - Tx) \leq (a + 2p + 1)F(Tx - x) \quad \forall x \in C. \tag{1.7}$$

Since C is convex, therefore $z = (1/2)T^2 x + (1/2)T^3 x$ is in C , and from the properties of the F -norm, we have

$$\begin{aligned} F(Tz - z) &\leq \frac{1}{2}F(Tz - T^2 x) + \frac{1}{2}F(Tz - T^3 x) \\ &\leq \frac{1}{2}\{aF(z - Tx) + bF(Tz - z) + cF(Tx - T^2 x) \\ &\quad + eF(Tx - Tz) + fF(z - T^2 x)\} \\ &\quad + \frac{1}{2}\{aF(z - T^2 x) + bF(Tz - z) + cF(T^3 x - T^2 x) \\ &\quad + eF(T^2 x - Tz) + fF(z - T^3 x)\}, \end{aligned}$$

4 Fixed Point Theory and Applications

$$\begin{aligned}
 F(z - Tx) &\leq \frac{1}{2}F(T^2x - Tx) + \frac{1}{2}F(T^3x - Tx) \\
 &\leq \frac{1}{2}F(Tx - x) + \frac{1}{2}(a + 2p + 1)F(Tx - x) = \left(1 + p + \frac{1}{2}a\right)F(Tx - x), \\
 F(z - T^2x) &\leq \frac{1}{2}F(T^3x - T^2x) \leq \frac{1}{2}F(Tx - x).
 \end{aligned}
 \tag{1.8}$$

Similarly,

$$\begin{aligned}
 F(z - T^3x) &\leq \frac{1}{2}F(Tx - x), \\
 F(Tx - Tz) &\leq \frac{1}{2}F(Tx - T^3x) + \frac{1}{2}F(Tx - T^4x) \\
 &\leq \frac{1}{2}(a + 2p + 1)F(Tx - x) + \frac{1}{2}\{F(Tx - T^2x) + F(T^2x - T^4x)\} \\
 &\leq \frac{1}{2}(a + 2p + 1)F(Tx - x) + \frac{1}{2}\{F(Tx - x) + (a + 2p + 1)F(Tx - x)\} \\
 &\leq \left(a + 2p + \frac{3}{2}\right)F(Tx - x), \\
 F(T^2x - Tz) &\leq \frac{1}{2}F(T^2x - T^3x) + \frac{1}{2}F(T^2x - T^4x) \leq \left(\frac{1}{2}a + p + 1\right)F(Tx - x).
 \end{aligned}
 \tag{1.9}$$

Thus

$$\begin{aligned}
 (1 - b)F(Tz - z) &\leq \frac{1}{2}\left\{a\left(1 + p + \frac{1}{2}a\right)F(Tx - x) + cF(Tx - x)\right. \\
 &\quad \left.+ e\left(a + 2p + \frac{3}{2}\right)F(Tx - x) + \frac{1}{2}fF(Tx - x)\right\} \\
 &\quad + \frac{1}{2}\left\{\frac{1}{2}aF(Tx - x) + cF(Tx - x) + \frac{1}{2}e(a + 2p + 1)F(Tx - x)\right. \\
 &\quad \left.+ \frac{1}{2}fF(Tx - x)\right\} = \left(\frac{3}{4}a + \frac{1}{4}a^2 + \frac{5}{4}ap + \frac{5}{2}p + \frac{3}{2}p^2\right)F(Tx - x).
 \end{aligned}
 \tag{1.10}$$

Thus

$$\begin{aligned}
 4(1 - p)F(z - Tz) &\leq (3a + a^2 + 5ap + 10p + 6p^2)F(Tx - x) \\
 &\leq (2p^2 - 5p + 4)F(Tx - x).
 \end{aligned}
 \tag{1.11}$$

Hence

$$\begin{aligned}
 F(z - Tz) &\leq \frac{26 - 22a - a^2}{8(a + 3)}F(Tx - x), \\
 F(Tz - z) &\leq \lambda F(Tx - x),
 \end{aligned}
 \tag{1.12}$$

where $\lambda = (26 - 22a - a^2)/8(a + 3)$. It is clear that $0 < \lambda < 1$.

Now let $i = \inf\{F(Tx - x) : x \in C\}$. Then there exists a point $x \in C$ such that $F(Tx - x) < i + \epsilon$ for $\epsilon > 0$.

Suppose $i > 0$. Then for $0 < \epsilon < (1 - \lambda)i/\lambda$ and $F(Tx - x) < i + \epsilon$, we have

$$F(Tz - z) \leq \lambda F(Tx - x) \leq \lambda(i + \epsilon) < i, \tag{1.13}$$

that is, $F(Tz - z) < i$, which is a contradiction with the definition of i . Hence $\inf\{F(Tx - x) : x \in C\} = 0$.

To prove that the infimum is attained is the easy part of the proof. Take the following system of sets: $K_n = \{x : F(x - Tx) \leq 1/2n(q + 1)\}$; $T(K_n)$ and $\overline{T(K_n)}$, where $n \in \mathbb{N}$, $q = (a + p)/(1 - a)$, and $\overline{T(K_n)}$ is the closure of $T(K_n)$. Then for any $x, y \in K_n$,

$$\begin{aligned} F(Tx - Ty) &\leq qF(Tx - x) + qF(Ty - y) \leq \frac{1}{n}, \\ F(x - y) &\leq (q + 1)F(Tx - x) + (q + 1)F(Ty - y) \leq \frac{1}{n}, \end{aligned} \tag{1.14}$$

that is, $\text{diam}(K_n) \leq 1/n$, $\text{diam}(T(K_n)) \leq 1/n$ and therefore, since $\text{diam}(T(K_n)) = \text{diam}(\overline{T(K_n)})$, we have $\text{diam}(\overline{T(K_n)}) \leq 1/n$. It is clear that $\{K_n\}$ and $\{\overline{T(K_n)}\}$ form monotone sequences of sets and from (1.5) we have $T(K_n) \subset K_n$. Suppose $y \in \overline{T(K_n)}$, then there exists $y' \in K_n$ such that $F(y - Ty') < \epsilon$ for $\epsilon > 0$ and

$$\begin{aligned} F(y - Ty) &\leq F(y - Ty') + F(Ty' - Ty) \\ &\leq F(y - Ty') + aF(y - y') + bF(y' - Ty') \\ &\quad + cF(Ty - y) + eF(y - Ty') + fF(y' - Ty). \end{aligned} \tag{1.15}$$

Hence

$$(1 - c)F(y - Ty) \leq (1 + a + e + f)\epsilon + (a + b)F(Ty' - y'). \tag{1.16}$$

Since $F(y' - Ty') \leq 1/2n(q + 1)$, then

$$F(y - Ty) \leq \frac{1 + a + e + f}{1 - c}\epsilon + \frac{a + b}{1 - c} \frac{1}{2n(q + 1)}. \tag{1.17}$$

Since $\epsilon > 0$ is arbitrary and $a + b + c \leq 1$, then $F(y - Ty) \leq 1/2n(q + 1)$ and we have $y \in K_n$. Hence $\overline{T(K_n)} \subset K_n$, too.

$\{\overline{T(K_n)}\}$ is a decreasing sequence of closed nonempty sets with $\text{diam}(\overline{T(K_n)}) \rightarrow 0$ as $n \rightarrow \infty$. Hence they have a nonempty intersection $\{x^*\}$ and T has a unique fixed point $Tx^* = x^*$. \square

COROLLARY 1.4. *Let C be a closed convex subset of a Banach space X and $T : C \rightarrow C$ a mapping that satisfies $\|Tx - Ty\| \leq a\|x - y\| + b\|Tx - x\| + c\|Ty - y\| + e\|Tx - y\| + f\|Ty - x\|$ for all $x, y \in C$ where $0 < a < 1$, $b \geq 0$, $c \geq 0$, $e \geq 0$, $f \geq 0$, and $a + b + c + e + f = 1$. Then T has a unique fixed point.*

COROLLARY 1.5 [1]. *Let C be a closed convex subset of a Banach space X and $T : C \rightarrow C$ a mapping that satisfies $\|Tx - Ty\| \leq a\|x - y\| + b\|Tx - x\| + c\|Ty - y\|$ for all $x, y \in C$, where $0 < a < 1$, $b \geq 0$, $c \geq 0$, and $a + b + c = 1$. Then T has a unique fixed point.*

6 Fixed Point Theory and Applications

COROLLARY 1.6. *Let C be a closed convex subset of a complete metrizable topological vector space X and $T : C \rightarrow C$ a mapping that satisfies $\|Tx - Ty\| \leq a\|x - y\| + b\|Tx - y\| + c\|Ty - x\|$ for all $x, y \in C$, where $0 < a < 1$, $b \geq 0$, $c \geq 0$, and $a + b + c = 1$. Then T has a unique fixed point.*

We now proceed to use the Mann iteration scheme [16] to approximate the fixed point of our mapping under consideration.

THEOREM 1.7. *Let C be a nonempty closed convex subset of a complete metrizable topological vector space X and let $T : C \rightarrow C$ be a mapping that satisfies $F(Tx - Ty) \leq aF(x - y) + bF(Tx - x) + cF(Ty - y) + eF(Tx - y) + fF(Ty - x)$ for all $x, y \in C$, where $0 < a < 1$, $b \geq 0$, $c \geq 0$, $e \geq 0$, $f \geq 0$, and $a + b + c + e + f = 1$. Suppose $\{x_n\}$ is a Mann iteration sequence defined by $x_{n+1} = (1 - \alpha_n)x_n + \alpha_nTx_n$, $x_0 \in C$, $n \geq 0$, where $\{\alpha_n\}$ satisfy $0 < \alpha_n \leq 1$ for all n , $\sum_0^\infty \alpha_n = \infty$. Assume $2c < c + b$, then $\{x_n\}$ converges to the unique fixed point of T .*

Proof. The fact that T has a unique fixed point is already shown in Theorem 1.3.

If $F(Tx - Ty) \leq aF(x - y) + bF(Tx - x) + cF(Ty - y) + eF(Tx - y) + fF(Ty - x)$, then

$$\begin{aligned} F(Tx - Ty) &\leq aF(x - y) + bF(Tx - x) + c\{F(Ty - Tx) + F(Tx - x) + F(x - y)\} \\ &\quad + e\{F(Tx - x) + F(x - y)\} + f\{F(Ty - Tx) + F(Tx - x)\}. \end{aligned} \quad (1.18)$$

After computation, we have $F(Tx - Ty) \leq ((a + c + e)/(1 - (c + f)))F(x - y) + ((b + c + e + f)/(1 - (c + f)))F(Tx - x)$. If $\delta = (a + c + e)/(1 - (c + f))$, then

$$F(Tx - Ty) \leq \delta F(x - y) + \frac{b + c + e + f}{1 - (c + f)} F(Tx - x). \quad (1.19)$$

Since by assumption $2c < b + c$, it is clear that $\delta < 1$.

Suppose p is a fixed point of T , then if $x = p$ and $y = x_n$, from (1.19), we obtain

$$\begin{aligned} F(Tx_n - p) &\leq \delta F(x_n - p), \\ F(x_{n+1} - p) &= F((1 - \alpha_n)x_n + \alpha_nTx_n - (1 - \alpha_n + \alpha_n)p) \\ &= F((1 - \alpha_n)(x_n - p) + \alpha_n(Tx_n - p)) \\ &\leq (1 - \alpha_n)F(x_n - p) + \alpha_nF(Tx_n - p) \\ &\leq (1 - \alpha_n(1 - \delta))F(x_n - p). \end{aligned} \quad (1.20)$$

Since $1 - \alpha_n(1 - \delta) < 1$ by the choice of α_n in the theorem, then $\{x_n\}$ converges to p . \square

Remarks 1.8. (1) Gregus [1] gave an example in which $a = 1$, C is closed convex and bounded but yet T does not have a fixed point. If $a = 1$, some form of boundedness must be assumed on C for T to have a fixed point, for example, see [7, 6]. The same is true if $a = 0$ (see [8, 9]).

(2) If (X, d) is a complete metric space and $a + b + c + e + f < 1$, it was shown in [17] that T as defined in (1.2) has a unique fixed point. However, if $a + b + c + e + f = 1$, Hardy

and Rogers [17] assumed that T is continuous and X is compact in order to prove the existence of fixed point for T as defined in (1.2). Goebel et al. [18] obtained the existence of fixed point for T as defined by (1.2) when $a + b + c + e + f = 1$. In which case, it was assumed that X is a uniformly convex Banach space, T is continuous and C is bounded, closed, and convex. In our result, T is not assumed to be continuous, X is assumed to be neither a compact nor a uniformly convex Banach space, and there is no boundedness assumption on C .

(3) Berinde [14] showed that the Ishikawa iteration sequence [16] of a class of quasi-contractive operators, called Zamfirescu operators, defined on a closed convex subset C of a Banach space X converges to the fixed point of T . The first author [19] showed that if X is a complete metrizable locally convex space, and C is closed and convex, then the Mann iteration sequence of the Zamfirescu operator T defined on C converges to the fixed point of T . In both cases, the sum of the constants is less than 1 while in Theorem 1.7, the sum is 1. In addition, X is generalized to a complete metrizable topological vector spaces. Can Theorem 1.7 still be proved without the assumption that $2c < a + b$?

References

- [1] M. Greguš Jr., "A fixed point theorem in Banach space," *Bollettino. Unione Matematica Italiana. A. Serie V*, vol. 17, no. 1, pp. 193–198, 1980.
- [2] P. P. Murthy, Y. J. Cho, and B. Fisher, "Common fixed points of Greguš type mappings," *Glasnik Matematički. Serija III*, vol. 30(50), no. 2, pp. 335–341, 1995.
- [3] R. N. Mukherjee and V. Verma, "A note on a fixed point theorem of Greguš," *Mathematica Japonica*, vol. 33, no. 5, pp. 745–749, 1988.
- [4] J. O. Olaleru, "A generalization of Greguš fixed point theorem," *Journal of Applied Sciences*, vol. 6, no. 15, pp. 3160–3163, 2006.
- [5] C. E. Chidume, "Geometric properties of Banach spaces and nonlinear iterations," Research Monograph, International Centre for Theoretical Physics, Trieste, Italy, in preparation.
- [6] A. Kaewcharoen and W. A. Kirk, "Nonexpansive mappings defined on unbounded domains," *Fixed Point Theory and Applications*, vol. 2006, Article ID 82080, 13 pages, 2006.
- [7] W. A. Kirk, "A fixed point theorem for mappings which do not increase distances," *The American Mathematical Monthly*, vol. 72, no. 9, pp. 1004–1006, 1965.
- [8] R. Kannan, "Some results on fixed points. III," *Fundamenta Mathematicae*, vol. 70, no. 2, pp. 169–177, 1971.
- [9] C. S. Wong, "On Kannan maps," *Proceedings of the American Mathematical Society*, vol. 47, no. 1, pp. 105–111, 1975.
- [10] S. K. Chatterjea, "Fixed-point theorems," *Comptes Rendus de l'Académie Bulgare des Sciences*, vol. 25, pp. 727–730, 1972.
- [11] J. O. Olaleru, "On weighted spaces without a fundamental sequence of bounded sets," *International Journal of Mathematics and Mathematical Sciences*, vol. 30, no. 8, pp. 449–457, 2002.
- [12] H. H. Schaefer and M. P. Wolff, *Topological Vector Spaces*, vol. 3 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 2nd edition, 1999.
- [13] N. Adasch, B. Ernst, and D. Keim, *Topological Vector Spaces*, vol. 639 of *Lecture Notes in Mathematics*, Springer, Berlin, 1978.
- [14] V. Berinde, "On the convergence of the Ishikawa iteration in the class of quasi contractive operators," *Acta Mathematica Universitatis Comenianae. New Series*, vol. 73, no. 1, pp. 119–126, 2004.

8 Fixed Point Theory and Applications

- [15] G. Köthe, *Topological Vector Spaces. I*, vol. 159 of *Die Grundlehren der mathematischen Wissenschaften*, Springer, New York, NY, USA, 1969.
- [16] B. E. Rhoades, "Comments on two fixed point iteration methods," *Journal of Mathematical Analysis and Applications*, vol. 56, no. 3, pp. 741–750, 1976.
- [17] G. E. Hardy and T. D. Rogers, "A generalization of a fixed point theorem of Reich," *Canadian Mathematical Bulletin*, vol. 16, pp. 201–206, 1973.
- [18] K. Goebel, W. A. Kirk, and T. N. Shimi, "A fixed point theorem in uniformly convex spaces," *Bollettino. Unione Matematica Italiana. Serie IV*, vol. 7, pp. 67–75, 1973.
- [19] J. O. Olaleru, "On the convergence of Mann iteration scheme in locally convex spaces," *Carpathian Journal of Mathematics*, vol. 22, no. 1-2, pp. 115–120, 2006.

J. O. Olaleru: Mathematics Department, University of Lagos, P.O. Box 31, Lagos, Nigeria
Email address: olaleru1@yahoo.co.uk

H. Akewe: Mathematics Department, University of Lagos, P.O. Box 31, Lagos, Nigeria