

Altering rainfall patterns through aerosol dispersion

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 J. Phys.: Conf. Ser. 852 012008

(<http://iopscience.iop.org/1742-6596/852/1/012008>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 197.210.200.118

This content was downloaded on 06/06/2017 at 17:43

Please note that [terms and conditions apply](#).

You may also be interested in:

[RAINFALL AT MOUNT HAMILTON \(1880-91\)](#)

[Experimental approach to validation of an analytical and numerical thermal analysis of a travelling wave tube](#)

W. Wiejak and A. Wymysowski

[Trends in the aerosol load properties over south eastern Italy](#)

J A G Orza and M R Perrone

[Formation of the CME Leading Edge Observed in the 2003 February 18Event](#)

Xing-Ming Bao, Hong-Qi Zhang and Jun Lin

[Retrieval of the aerosol optical thickness from UV global irradiance measurements](#)

M J Costa, V Salgueiro, D Bortoli et al.

[Heating of heavy oil reservoirs using high-temperature heat agent recirculation](#)

V B Osnos, V V Kuneevsky, V M Larionov et al.

[Analysis of atmospheric vertical profiles in the presence of desert dust aerosols](#)

M J Costa, M A Obregón, S Pereira et al.

[Detection of fires and explosions](#)

B. Ray

[Hygroscopic Properties of Atmospheric Aerosol Measured with an HTDMA in an Urban Background Site in Madrid](#)

E Alonso-Blanco, F J Gómez-Moreno, M Becerril et al.

Altering rainfall patterns through aerosol dispersion

M E Emeteri¹, M Bakeko,¹ L Onyechekwa¹ and W Ayara¹

¹Physics Department, Covenant University, P.M.B. 1023, Ota-Ogun State, Nigeria

E-mail : moses.emeteri@covenantuniversity.edu.ng

Abstract. The possibility of recirculation mechanism on rainfall patterns is salient for sustenance of the human race through agricultural produce. The peculiarity of the lower atmosphere of south west region of Nigeria was explored using theoretical and experimental approach. In the theoretical approach, the reconstruction of 1D model as an extraction from the 3D aerosol dispersion model was used to examine the physics of the recirculation theory. The experimental approach which consists of obtaining dataset from ground instruments was used to provide on-site guide for developing the new recirculation theories. The data set was obtained from the Davis weather station, Nigeria Meteorological agency and Multi-angle Imaging Spectro-radiometer (MISR). We looked at the main drivers of recirculation and propounded that recirculation is a complex process which triggers a reordering of the mixing layer- a key factor for initiating the type of rainfall in this region.

1. Introduction

The role of aerosols in understanding the climatic system is enormous because it controls verse processes in the atmosphere like atmospheric circulation [1], rainfall pattern [2] cloud formation [3]. Its direct and indirect influence on the global climate can be very complicating when understudying the comparative role of aerosols to global energy balance [4]. Atmospheric aerosols have been reported to form heterogeneous or homogenous layers in the atmosphere [5]. Models are propounded to amplify the usefulness and exactness of the measuring instruments. Sadly, these model needs change in a few zones e.g. sufficient estimation of airborne scattering and statement, rates of new molecule arrangement and their reliance on controlling factors e.g. development and maturing. The aerosols dispersion and deposition technique requires a systemic evaluation of aerosols not only at ground level but also in the planetary boundary layer (PBL) to determine the aerosol transport mechanism-in relation to weather formation at the PBL [5, 6]. The planetary boundary layer contains a large variety of aerosols from anthropogenic and natural sources [1]. The planetary boundary layer (PBL) comprises of sub-layers e.g. mixed layer, surface layer, stable layer and residual layer. The surface layer is closest to the earth i.e. 10 percent of the height of the PBL. Turbulent stresses are relatively constant. The mixed layer or convective layer are directly above the surface layer and are characterized by convective motion and significant mixing of air. It makes up the entire layer of the PBL above the surface layer - during day and reduces drastically during the night. The stable layer is located above the mixed layer. It is the entry point where air from above the PBL entrains into the mixed layer. This stable layer restricts turbulence, prevents frictional influences from reaching above the PBL and prevents the continued upward motion of thermals. The residual layer is majorly residuals of the mixed layer and is not influenced by turbulent stresses. The PBL are influenced by environment forces like differential heating, energy budgets, moisture, diurnal variations, buoyancy, shear, and roughness length. The PBL takes out aerosol retention by means of regular components which works



either as wet deposition, dry deposition or chemical reactions. These procedures can happen exclusively or partially. Layering of the PBL is subject to its temperature. All the more for all intents and purposes, the review site that is featured in Emeter et al. [6] transverses the beach front area (in the south of Nigeria) to the sub-Sahel locale (in the north of Nigeria). Significantly, the wind transport instrument starts turbulent weight on the PBL. For instance, amid harmattan, the wind transports vast amounts of particulates from the Sahara Desert in the north toward the south. Likewise, aerosols due to bush burning, automobile activities in the south increase the aerosols content over the study site [6]. Amid the non-harmattan, the West African Sub area is washed by the sticky SW Monsoon winds from the Atlantic Ocean [7].

The recirculation theory helps us to understand the anomalies in both the aerosols dispersion and deposition points of the atmospheric field settings. The mysteries of the recirculation zones in the mixing layer of the lower atmosphere are yet to be explained i.e. comparing the pre-circulation and post circulation. The turbulent boundary layers formed during recirculation are somewhat complex when explaining the minute changes that triggers continuous recirculation [8, 9]. In this paper, we understudied a 3D dispersion model which was developed from the aerosols dispersion dynamics around a cement factory. We looked at the main drivers of recirculation and propounded that recirculation is a complex process which triggers a reordering of the mixing layer- a key factor for initiating the type of rainfall in this region.

2. Methodology

The Davis weather station used for this study was installed at Covenant University, Ota Nigeria. Ota is on 6.21°N and 3.21°E. The weather station was used to obtain the wind properties i.e. peculiar to the region. The Meteorological data for rainfall was obtained from the Nigerian Meteorological Agency for five States in the south-west i.e. Osogbo-Osun, Ilorin-Kwara, Ibadan-Oyo, Abeokuta-Ogun, Ondo-Ondo and Ikeja-Lagos. The main objective of the theoretical approach is to derive recirculation equations from micro scale. The theoretical approach to this studies can be obtained from the aerosol dispersion model [5, 6]. The mathematical representation of type 3 is given as

$$\frac{\partial C}{\partial t} + V_x \frac{\partial C}{\partial x} - V_z \frac{\partial C}{\partial z} - V_y \frac{\partial C}{\partial y} = \frac{\partial}{\partial z} \left(K_z \frac{\partial C}{\partial z} \right) + \frac{\partial}{\partial y} \left(K_y \frac{\partial C}{\partial y} \right) + \frac{\partial}{\partial z} \left(K_{z2} \frac{\partial C}{\partial z} \right) + \frac{\partial}{\partial y} \left(K_{y2} \frac{\partial C}{\partial y} \right) - B + D \quad (1)$$

Here D represents the transport equations for recirculation; B is the pressure gradient force.

$$D = \frac{\partial(V_x \epsilon)}{\partial x} = \frac{\partial}{\partial x} \left(\frac{V_t}{\sigma_\epsilon} \frac{\partial \epsilon}{\partial x} \right) - k \quad (2)$$

$$B = dP. dA \quad (3)$$

Here V_x is the mean velocity in x direction, P is the pressure, A is the surface area, V_t is the kinematic viscosity, ϵ is the turbulence kinetic energy, k is the dissipation rate, σ_ϵ turbulent diffusivity coefficient.

$$\frac{\partial C}{\partial t} + V_x \frac{\partial C}{\partial x} = V_x \frac{\partial C}{\partial x} + \frac{\partial}{\partial x} \left(\frac{V_t}{\sigma_\epsilon} \frac{\partial \epsilon}{\partial x} \right) - k \quad (4)$$

Equation 7 has important implications for precipitation formation via the recirculation region. The recirculation region initiates the nucleation scavenging which grows the precipitation size. The ability of aerosol particles to act as a cloud condensation nuclei (CCN) depends on its size [10]. The solution of equation (4) is given as

$$\epsilon = \frac{\sigma_\epsilon}{V_t} (-V_x(kt + C - x) - PAx) \quad (5)$$

The solution i.e. equation (5) shows that the updraft term $xV_x \frac{\sigma_\epsilon}{V_t}$ was able to initiate the nucleation scavenging. Hence, large scale recirculation may influence precipitation over an area provided that

other terms in equation (5) are quite low. This assumption was verified by the rainfall and aerosol optical depth over the research site.

3. Results and Discussion

In this section we tried verifying how intense is the relationship between the aerosol content of an area and the rainfall pattern. The data duration is thirteen years. First we examine the wind speed data (Figures 1(a) and 1(b)) from the Davis weather station to determine the various expected possibilities in the recirculation zones.

The wind data set for January, 2013 was used to estimate the recirculation zone in 1D where $x=z$. More specifically, the recirculation zone occurs more frequently along the SE, S, ESE, ENE, E, SSW and SSE as shown in Figure 1(b). If we assume that all the negative terms in equation 8 are minute, then the turbulence kinetic energy which defines the sustenance of the recirculation zone can be written as

$$\epsilon \approx xV_x \frac{\sigma_\epsilon}{V_t} \quad (6)$$

where

$$V_x(\theta) = \begin{cases} V\sin\theta & \text{if } 0 < x < 480, \quad 0 < \theta < \pi/16 \\ V\cos\theta & \text{if } 480 \leq x < 960, \quad \pi/16 < \theta < \pi/8 \\ V(\sin\theta + \cos\theta) & \text{if } 960 \leq x < 1440, \quad \pi/8 < \theta < \pi/4 \\ V\sin^2\theta & \text{if } 1440 \leq x < 1920, \quad \pi/4 < \theta < \pi/2 \\ V\cos^2\theta & \text{if } 1920 \leq x < 2400, \quad \pi/2 < \theta < \pi \end{cases}$$

$\frac{\sigma_\epsilon}{V_t} = 0.01$, and $V = 7.8\text{m/s}$ i.e. the highest speed with an equivalently high frequency. Let x be the altitude of the lower atmosphere where altitude 1440 m to 2400 m represents the free troposphere. The recirculation activity was theoretically represented in Figure 2.
















Figure 1: Magnitude of wind speed within different directions

Figure 1. (a) Magnitude of wind speed within different directions and **(b)** Frequency of wind directions

The 3D representation of the recirculation parameters illustrated in equation (9) where the maximum turbulence kinetic energy over the research area-which is about 261 J/Kg and occurs between 1000 m to 1500 m altitude (see Fig. 3). The divergence of recirculation features as seen by the red dotted lines is evidence that the recirculation zones are unpredictably dynamic. Hence, it possesses a verse ability to influence rainfall over an area. We understudy the rainfall pattern between the years 2000 to 2012 over the research area. A comparative analysis between the rainfall pattern and the aerosol optical depth was carried to further explain the dynamic nature of the recirculation zones. This study provides the hint into understanding the tropospheric anomaly noticed by Lee et al. [11]. Lee et al. [11] proposed that via statistical analysis that biomass burning aerosol (BBA) affects liquid clouds- leading to an increase in cloud droplet number concentrations and a decrease of droplet

effective radii. In the long run, this process leads to geographical rearrangements of rainfall patterns. The theory of recirculation as explain in this paper sheds more light on the possibility of rainfall suppression and promotion [12, 13]. The recirculation activity can influence the normal deceleration (in form of normalization, accelerating or decreasing) of auto-conversion processes in the lower atmosphere [14].








Figure 2. Rainfall pattern within the years 2000 to 2012




Figure 3. (a) AOD pattern within the years 2000 to 2012 for Lagos and (b) AOD pattern within the years 2000 to 2012 for Ilorin.

4. Conclusion

The recirculation zone is comprised the negative recirculation which supports the downdraft activities and positive recirculation which supports the updraft activities within the lower atmospheric setting. The negative recirculation has no influence on the rainfall pattern of an area-as seen in Osogbo-Osun. The positive recirculation has significant influence on the rainfall pattern of an area-as seen in Ikeja-Lagos. The maximum turbulence kinetic energy over the research area is about 261 J/kg and occurs between 1000 m to 1500m altitude. This altitude shows the role of recirculation in cloud formation and rainfall. Hence, the meteorological data set for the six locations shows that Ondo and Ikeja-Lagos has the highest value of turbulence kinetic energy. Also, we discover that the turbulence kinetic energy of negative recirculation is low i.e. compared with the turbulence kinetic energy of positive influence. The rainfall pattern of 2001 seems to be uniquely uniform with a clear evidence of uniform recirculation activity throughout the six locations. It was shown that recirculation activity influences the nature of clouds. The glaring results are the increase in cloud droplet number concentrations and a

decrease of droplet effective radii. This result is possible because nucleation scavenging triggers rainfall both at the long or short term as shown in the results.

Acknowledgement

The authors acknowledge the Nigeria Meteorological Agency, NASA and Covenant University for allowing us to use their data set.

References

- [1] Andreas Bott 1999 A numerical model of the cloud-topped planetary boundary-layer: chemistry in marine stratus and the effects on aerosol particles *Atmospheric Environment* **33**(12)1921
- [2] Rotstayn L D and Lohmann U 2002 Tropical rainfall trends and the indirect aerosol effect *J. Clim* **15**(15) 2103
- [3] Allen R J, Sherwood S C, 2010 The aerosol-cloud semi-direct effect and land-sea temperature contrast in a GCM *Geophys. Res. Lett.* **37** L07702
- [4] Mao K B, Ma Y, Xia L, Wendy Y C *et al.* 2014 Global aerosol change in the last decade: An analysis based on MODIS data *Atmospheric Environment* **94** 680
- [5] Emeteri M E, Akinyemi M L and Akin-Ojo O 2015 Aerosol Optical Depth Trends over Different Regions of Nigeria: Thirteen years Analysis *Modern Applied Science* **9**(9) 267
- [6] Emeteri M E , Akinyemi M L , Uno U E 2015 Computational analysis of aerosol dispersion trends from cement factory *Proceedings 2015 International Conference on Space Science & Communication* p 288
- [7] Okpani P E 2009 Determination of aerosol mass loading in ilorin using remote sensing technique *The Pacific Journal of Science and Technology* **10**(2) 5
- [8] Sadek R, Soulhac L, Brocheton F, Buisson E 2011 Evaluation of wind field and dispersion models in complex terrain, 14th HARMO conference pp 1-23
- [9] Kim H G and Patel V C 200 Test of turbulence models for wind flow over terrain with separation and recirculation *Boundary-Layer Meteorology* **94** 5
- [10] Westervelt D M, Pierce J R and Adams P J 2014 Analysis of feedbacks between nucleation rate, survival probability and cloud condensation nuclei formation *Atmos. Chem. Phys.* **14** 5577
- [11] Lee D, Sud Y C, Oreopoulos L *et al.* 2014 Modeling the influences of aerosols on pre-monsoon circulation and rainfall over Southeast Asia *Atmos. Chem. Phys.* **14** 6853
- [12] Iacobellis S F and Somerville R C J 2006 Evaluating parameterizations of the autoconversion process using a single-column model and atmospheric radiation measurement program measurements *J. Geophys. Res.* **111** D02203
- [13] Diem J E 2013 Response to comments on Influences of the Bermuda High and atmospheric moistening on changes in summer rainfall in the Atlanta, Georgia region, the United States *International Journal of Climatology* **33** 3086
- [14] Weber T and J Quaas 2012 Incorporating the subgrid-scale variability of clouds in the autoconversion parameterization using a PDF-scheme *J. Adv. Model. Earth Syst.* **4** M11003